

COMPACTLY GENERATED STACKS: A CARTESIAN-CLOSED THEORY OF TOPOLOGICAL STACKS

DAVID CARCHEDI

ABSTRACT. A convenient 2-category of topological stacks is constructed which is both complete and Cartesian closed. This 2-category, called the 2-category of compactly generated stacks, is the analogue of classical topological stacks, but for a different Grothendieck topology. In fact, there is an equivalence of 2-categories between compactly generated stacks and those classical topological stacks which admit locally compact atlases. Compactly generated stacks are also equivalent to a bicategory of topological groupoids and principal bundles, just as in the classical case. If a classical topological stack and a compactly generated stack have a presentation by the same topological groupoid, then they restrict to the same stack over locally compact Hausdorff spaces and are homotopy equivalent.

CONTENTS

1. Introduction	2
1.1. Why are Compactly Generated Hausdorff Spaces Cartesian Closed?	3
1.2. Organization and Main Results	4
2. Topological Stacks	7
2.1. Fibrant Topological Groupoids	7
2.2. The Homotopy Type of a Topological Stack	8
2.3. Homotopical Stacks and Mapping Stacks	8
3. The Compactly Generated Grothendieck Topology	11
3.1. The Compactly Generated Grothendieck Topology	11
3.2. Stacks for the Compactly Generated Grothendieck Topology	18
4. Compactly Generated Stacks	19
4.1. Compactly Generated Stacks	19
4.2. Mapping Stacks of Compactly Generated Stacks	23
4.3. Homotopy Type of Compactly Generated Stacks	25
4.4. Comparison with Topological Stacks	26
Appendix A. Stacks	28
Appendix B. Topological Groupoids and Topological Stacks	31
B.1. Definitions and Some Examples	31
B.2. Principal Bundles	32
B.3. Bicategories of Topological Groupoids	34
B.4. Topological Stacks	36
B.5. Representable Maps of Topological Stacks	38
References	38

1. INTRODUCTION

The aim of this paper is to introduce the 2-category of compactly generated stacks. Compactly generated stacks are “essentially the same” as topological stacks, however, their associated 2-category is Cartesian-closed and complete, whereas the 2-category of topological stacks appears to enjoy neither of these properties. In this paper, we show that these categorical shortcomings can be overcome by refining the open cover Grothendieck topology to take into account compact generation.

The study of the mapping stack between topological stacks has been done in many different settings. The special case of differentiable maps between orbifolds has been studied in [3], and is restricted to the case where the domain orbifold is compact. André Haefliger has studied the case of smooth maps between étale Lie groupoids (which correspond to differentiable stacks with an étale atlas) in [8]. In [9], Ernesto Lupercio and Bernardo Uribe showed that the free loop stack (the stack of maps from S^1 to the stack) of an arbitrary topological stack is again a topological stack. In [20], Behrang Noohi addressed the general case of maps between topological stacks. He showed that under a certain compactness condition on the domain stack, the stack of maps between two topological stacks is a topological stack and if this compactness condition is replaced with a local compactness condition, the mapping stack is “not very far” from being topological.

In order to obtain a Cartesian-closed 2-category of topological stacks, we first restrict to stacks over a Cartesian-closed subcategory of the category $\mathbf{TOP}$ of all topological spaces. For instance, all of the results of [20] about mapping stacks are about stacks over the category of compactly generated spaces with respect to the open cover Grothendieck topology. We choose to work over the category of compactly generated Hausdorff spaces since, in addition to being Cartesian-closed, every compact Hausdorff space is locally compact Hausdorff, which is crucial in defining the compactly generated Grothendieck topology.

There are several equivalent ways of describing compactly generated stacks. The description that substantiates most clearly the name “compactly generated” is the description in terms of topological groupoids and principal bundles. Recall that the 2-category of topological stacks is equivalent to the bicategory of topological groupoids and principal bundles. Classically, if X is a topological space and $\mathcal{G}$ is a topological groupoid, the map π of a (left) principal $\mathcal{G}$ -bundle P over X

$$\begin{array}{ccc} \mathcal{G}_1 & \xrightarrow{\mathcal{C}} & P \\ \downarrow & \swarrow \mu & \downarrow \pi \\ \mathcal{G}_0 & & X \end{array}$$

must admit local sections. The notion of a principal bundle makes sense without this requirement; this local triviality condition is precisely where the open cover Grothendieck topology manifests itself. Let us call an object that would otherwise be a principal bundle, if not for this bundle map admitting local sections, simply a principal bundle, and call one which admits local sections an *ordinary* principal bundle. A compactly generated principal bundle is then a principal bundle whose restriction to any compact subset of X is an ordinary principal bundle. With this notion of compactly generated principal bundles, one can define a bicategory of

topological groupoids in an obvious way. This bicategory is equivalent to compactly generated stacks.

There is another simple way of defining compactly generated stacks. Given any stack $\mathcal{X}$ over the category of compactly generated Hausdorff spaces, it can be restricted to the category of compact Hausdorff spaces $\mathbb{CH}$. This produces a 2-functor

$$j^* : \mathfrak{TSt} \rightarrow St(\mathbb{CH})$$

from the 2-category of topological stacks to the 2-category of stacks over compact Hausdorff spaces. Compactly generated stacks are (equivalent to) the essential image of this 2-functor.

Finally, the simplest description of compactly generated stacks is that compactly generated stacks are classical topological stacks (over compactly generated Hausdorff spaces) which admit a locally compact atlas. In this description, the mapping stack of two spaces is usually not a space, but a stack!

For technical reasons, neither of the three previous concepts of compactly generated stacks are put forth as the definition. Instead, a Grothendieck topology $\mathcal{CG}$ is introduced on the category $\mathbb{CGH}$ of compactly generated Hausdorff spaces which takes into account the compact generation of this category. It is in fact the Grothendieck topology induced by geometrically embedding the topos $Sh(\mathbb{CH})$ of sheaves over $\mathbb{CH}$ into the topos of presheaves $Set^{\mathbb{CGH}^{op}}$. Compactly generated stacks are defined to be presentable stacks (see Definition B.16) with respect to this Grothendieck topology. The equivalence of all four notions of compactly generated stacks is shown in Section 4.1.

1.1. Why are Compactly Generated Hausdorff Spaces Cartesian Closed?

In order to obtain a Cartesian closed 2-category of topological stacks, we start with a Cartesian closed category of topological spaces. We choose to work with the aforementioned category $\mathbb{CGH}$ of compactly generated Hausdorff spaces (also known as Kelley spaces).

Definition 1.1. A topological space X is **compactly generated** if it has the final topology with respect to all maps into X with compact Hausdorff domain. When X is Hausdorff, this is equivalent to saying that a subset A of X is open if and only if its intersection with every compact subset of X is open.

The inclusion $\mathbb{CGH} \hookrightarrow \mathbb{HAUS}$ of the category of compactly generated Hausdorff spaces into the category of Hausdorff spaces admits a right-adjoint, called the Kelley functor, which replaces the topology of a space X with the final topology with respect to all maps into X with compact Hausdorff domain. Limits in $\mathbb{CGH}$ are computed by first computing the limit in $\mathbb{HAUS}$, and then applying the Kelley functor (in this way the compactly generated product topology differs from the ordinary product topology). In particular, $\mathbb{CGH}$ is a complete category.

Although the fact that this category is Cartesian-closed is a classical result (see: [24]), we will recall briefly the key reasons why this is true in order to gain insight into how one could construct a Cartesian-closed theory of topological stacks.

- (1) In $\mathbb{TOP}$, if K is compact Hausdorff, then for any space X , the space of maps endowed with the compact-open topology serves as an exponential object $\mathbb{Map}(K, X)$.

- (2) A Hausdorff space Y is compactly generated if and only if it is the colimit of all its compact subsets:

$$Y = \varinjlim_{K_\alpha \hookrightarrow Y} K_\alpha.$$

- i) For a fixed Y , for all X , $\mathbb{M}\mathrm{ap}(K_\alpha, X)$ exists for each compact subset K_α of Y .
- ii) $\mathbb{C}\mathbb{G}\mathbb{H}$ has all limits

So by general properties of limits and colimits, the space

$$\mathbb{M}\mathrm{ap}(Y, X) := \varinjlim_{K_\alpha \hookrightarrow Y} \mathbb{M}\mathrm{ap}(K_\alpha, X)$$

is a well defined exponential object (with the correct universal property).

The story starts the same for topological stacks:

Let Y be as above and let $\mathcal{X}$ be a topological stack. Then $\mathbb{M}\mathrm{ap}(K_\alpha, \mathcal{X})$ is a topological stack for each compact subset $K_\alpha \subset Y$ (see [20]).

One might therefore be tempted to claim:

$$\mathbb{M}\mathrm{ap}(Y, \mathcal{X}) := \varprojlim_{K_\alpha \hookrightarrow Y} \mathbb{M}\mathrm{ap}(K_\alpha, \mathcal{X})$$

But, there are some problems with this. First of all, this weak 2-limit may not exist as a topological stack, since topological stacks are only known to have finite weak limits. There is also a more technical problem related to the fact that the Yoneda embedding does not preserve colimits (see Section 3 for details). The main task of this paper is to show that both of these difficulties can be surmounted by using a more suitable choice of Grothendieck topology on the category of compactly generated Hausdorff spaces. The resulting 2-category of presentable stacks with respect to this topology will be the 2-category of compactly generated stacks and turn out to be both Cartesian-closed and complete.

1.2. Organization and Main Results. Section 2 is a review of some recent developments in topological groupoids and topological stacks, including some results of David Gepner and André Henriques in [7] which are crucial for the proof of the completeness and Cartesian-closedness of compactly generated stacks. In this section, we also extend Behrang Noohi's results to show that the mapping stack of two topological stacks is “nearly topological” if the domain stack admits an atlas by a CW-complex.

Section 3 details the construction of the compactly generated Grothendieck topology $\mathcal{C}\mathcal{G}$ on the category of compactly generated Hausdorff spaces $\mathbb{C}\mathbb{G}\mathbb{H}$. This is the Grothendieck topology whose associated presentable stacks are precisely compactly generated stacks. Many properties of the associated categories of sheaves and stacks are derived.

Section 4 is dedicated to compactly generated stacks. In Section 4.1, it is shown that compactly generated stacks are equivalent to two bicategories of topological groupoids. Also, it is shown that these are in turn equivalent to the restriction of topological stacks to compact Hausdorff spaces. Finally, it is shown that compactly generated stacks are equivalent to ordinary topological stacks (over compactly generated Hausdorff spaces) which admit a locally compact atlas.

Section 4.1 also contains one of the main results of the paper:

Theorem 1.1. *The 2-category of compactly generated stacks is closed under arbitrary small weak limits.*

(See Corollary 4.2).

Section 4.2 is dedicated to the proof of *the* main result of the paper:

Theorem 1.2. *If $\mathcal{X}$ and $\mathcal{Y}$ are arbitrary compactly generated stacks, then $\mathbb{M}\mathrm{ap}(\mathcal{Y}, \mathcal{X})$ is a compactly generated stack.*

(See Theorem 4.8).

This of course proves that classical topological stacks (over compactly generated Hausdorff spaces) which admit a locally compact atlas form a Cartesian-closed complete 2-category.

Finally, in section 4.4, there is a series of results showing how compactly generated stacks are “essentially the same” as topological stacks.

For instance, the following proposition is proven:

Proposition 1.1. *Let $\mathcal{X}$ be a topological stack and let $St_{\mathcal{CG}}(\mathcal{X})$ be its $\mathcal{CG}$ -stackification. Then the canonical map*

$$\mathcal{X} \rightarrow St_{\mathcal{CG}}(\mathcal{X})$$

induces an equivalence of groupoids

$$\mathcal{X}(Y) \rightarrow St_{\mathcal{CG}}\mathcal{X}(Y)$$

for all locally compact Hausdorff spaces Y . Moreover, this map is a universal weak equivalence.

(See Proposition 4.2).

It is also shown that when $\mathcal{X}$ and $\mathcal{Y}$ are topological stacks and $\mathcal{Y}$ satisfies the local-compactness condition necessary in [20] for $\mathbb{M}\mathrm{ap}(\mathcal{Y}, \mathcal{X})$ to be “nearly topological”, then:

Theorem 1.3. *There is a universal weak equivalence*

$$\mathbb{M}\mathrm{ap}(\mathcal{Y}, \mathcal{X}) \xrightarrow{\sim} \mathbb{M}\mathrm{ap}(St_{\mathcal{CG}}(\mathcal{Y}), St_{\mathcal{CG}}(\mathcal{X})),$$

and these two stacks agree over locally compact paracompact Hausdorff spaces.

(See Theorem 4.10).

Finally, a concrete description of a topological groupoid presentation for the mapping stack of two compactly generated stacks is given. Using this, the previous result is extended to the case where $\mathcal{Y}$ admits a CW-complex as an atlas.

Acknowledgment: I would like to thank my adviser Ieke Moerdijk for his guidance and patience; it was due to him that I came to consider the question of the presentability of the mapping stack of two topological stacks, starting first through the language of topological groupoids and étendues. I would also like to thank Marius Crainic for sparking my interest, in the first place, in topological groupoids and Lie groupoids in the form of a very interesting seminar as part of an MRI masterclass. I would very much like to thank André Henriques as well for countless hours of useful mathematical discussions and suggestions. I would like to thank André Haefliger for taking the time to explain to me his work on maps between

étale Lie groupoids during a visit to Genève. I would also like to thank Behrang Noohi for his helpful e-mail correspondence. This paper was written during my (current) studies as a Ph.D. student at Utrecht University.

2. TOPOLOGICAL STACKS

A review of the basics of topological groupoids and topological stacks including many notational conventions used in this section can be found in Appendix B.

2.1. Fibrant Topological Groupoids. The notion of fibrant topological groupoids was introduced in [7]. Roughly speaking, fibrant topological groupoids are topological groupoids which “in the eyes of paracompact spaces are stacks.” The fact that every topological groupoid is Morita equivalent to a fibrant one is essential to the existence of arbitrary weak limits of compactly generated stacks. Since this concept is relatively new, in this subsection, we summarize the basic facts about fibrant topological groupoids. All details may be found in [7].

Definition 2.1. The **classifying space** of a topological groupoid $\mathcal{G}$ is the fat geometric realization of its simplicial nerve (regarded as a simplicial space) and is denoted by $\|\mathcal{G}\|$.

For any topological groupoid $\mathcal{G}$, the classifying space of its translation groupoid $\|\mathbb{E}\mathcal{G}\|$ (see definition B.8) admits the structure of a principal $\mathcal{G}$ -bundle over the classifying space $\|\mathcal{G}\|$.

Definition 2.2. [7] Let $\mathcal{G}$ be a topological groupoid. A principal $\mathcal{G}$ -bundle E over a space B is **universal** if every principal $\mathcal{G}$ -bundle P over a paracompact base X admits a $\mathcal{G}$ -bundle map

$$\begin{array}{ccc} P & \longrightarrow & E \\ \downarrow & & \downarrow \\ X & \longrightarrow & B, \end{array}$$

unique up to homotopy.

Lemma 2.1. [7] *The principal $\mathcal{G}$ -bundle*

$$\begin{array}{ccccc} \mathcal{G}_1 & \xrightarrow{\mathcal{G}} & \|\mathbb{E}\mathcal{G}\| \\ \downarrow & \searrow \mu & \downarrow \pi \\ \mathcal{G}_0 & & \|\mathcal{G}\| \end{array}$$

is universal.

Definition 2.3. [7] A topological groupoid $\mathcal{G}$ is **fibrant** if the unit principal $\mathcal{G}$ -bundle is universal.

Definition 2.4. [7] The **fibrant replacement** of a topological groupoid $\mathcal{G}$ is the gauge groupoid of the universal principal $\mathcal{G}$ -bundle $\|\mathbb{E}\mathcal{G}\|$, denoted $Fib(\mathcal{G})$.

Remark. If $\mathcal{G}$ is compactly generated Hausdorff, then so is $Fib(\mathcal{G})$.

Lemma 2.2. [7] *$Fib(\mathcal{G})$ is fibrant.*

Lemma 2.3. [7] *There is a canonical groupoid homomorphism*

$$\xi_{\mathcal{G}} : \mathcal{G} \rightarrow Fib(\mathcal{G})$$

which is a Morita equivalence for all topological groupoids $\mathcal{G}$.

The following theorem will be of importance later:

Theorem 2.4. [7] *Let $\mathcal{G}$ be a fibrant topological groupoid. Then for any topological groupoid $\mathcal{H}$ with paracompact Hausdorff object space $\mathcal{H}_0$, there is an equivalence of groupoids*

$$\mathrm{Hom}_{\mathfrak{T}\mathfrak{G}\mathfrak{t}}([\mathcal{H}], [\mathcal{G}]) \cong \mathrm{Hom}_{\mathrm{TOP}Gpd}(\mathcal{H}, \mathcal{G})$$

natural in $\mathcal{H}$. In particular, the restriction of $\tilde{y}_{\mathcal{G}}$ to the full-subcategory of paracompact Hausdorff spaces agrees with the restriction of $[\mathcal{G}]$.

(See definition B.17 for notation.)

2.2. The Homotopy Type of a Topological Stack. The weak homotopy type of a stack $\mathcal{X}$ over TOP with respect to the open cover topology is the homotopy type of a terminal object in the homotopy category of CW-complexes over $\mathcal{X}$. Such a weak homotopy type always exists. For details, see [7]. In the case that $\mathcal{X}$ is a topological stack, we have a concrete description of the weak homotopy type.

Proposition 2.1. [7] *If $\mathcal{X}$ is a topological stack, the **weak homotopy type** of $\mathcal{X}$ is the weak homotopy type of $\|\mathcal{G}\|$ for any topological groupoid $\mathcal{G}$ for which $\mathcal{X} \cong [\mathcal{G}]$.*

The above proposition is not new in that sense that it has been the working definition of the homotopy type of a topological stack for quite some time. For instance, this is the definition of the homotopy type of an orbifold given in [1].

As a corollary, we obtain the following:

Corollary 2.1. *If $\mathcal{G} \rightarrow \mathcal{H}$ is a Morita equivalence, the induced map $\|\mathcal{G}\| \rightarrow \|\mathcal{H}\|$ is a weak homotopy equivalence.*

This is a classical result. For instance, it is proven for the case of étale topological groupoids in [12] and [13].¹

Definition 2.5. A map $f : X \rightarrow Y$ of spaces is a **universal weak equivalence** if for any map $T \rightarrow Y$, the induced map $T \times_Y X \rightarrow T$ is a weak homotopy equivalence.

Remark. The property of being a universal weak equivalence is stable under restriction and local on the target.

It follows from Lemma 2.3 that given $\mathcal{X} \cong [\mathcal{G}]$, there exists an atlas

$$\varphi : \|\mathcal{G}\| \rightarrow \mathcal{X}.$$

Theorem 2.5. [19] *The atlas $\varphi : \|\mathcal{G}\| \rightarrow \mathcal{X}$ is a universal weak equivalence.*

2.3. Homotopical Stacks and Mapping Stacks. Stacks on TOP (with respect to the open cover topology) come in many different flavours. Of particular importance of course are topological stacks- those stacks coming from topological groupoids. However, this class of stacks seems to be too restrictive since many natural stacks, for instance the stack of maps between two topological stacks, appear to not be topological.

A topological stack is a stack $\mathcal{X}$ which admits an epimorphism $X \rightarrow \mathcal{X}$ from a topological space X . This implies:

¹I would like to thank Ieke Moerdijk for explaining to me how to extend his method of proof to any topological groupoid whose object and arrow spaces have a basis of contractible open sets.

- i) Any map $T \rightarrow \mathcal{X}$ from a topological space is representable (equivalently, the diagonal $\Delta : \mathcal{X} \rightarrow \mathcal{X} \times \mathcal{X}$ is representable) [18].
- ii) If $T \rightarrow \mathcal{X}$ is a continuous map, then the induced map $T \times_{\mathcal{X}} X \rightarrow T$ admits local sections (i.e. is an epimorphism in $\mathbf{TOP}$).

If the second condition is slightly weakened, the result is a stack which is “nearly topological”.

Definition 2.6. [19] A **paratopological** stack is a stack $\mathcal{X}$ on $\mathbf{TOP}$ (with respect to the open cover topology), satisfying condition i) above, and satisfying condition ii) for all maps $T \rightarrow \mathcal{X}$ from a paracompact space.

Definition 2.7. [19] A **pseudotopological** stack is a stack $\mathcal{X}$ on $\mathbf{TOP}$ (with respect to the open cover topology), satisfying condition i) above, and satisfying condition ii) for all maps $T \rightarrow \mathcal{X}$ from a CW-complex.

Paratopological and pseudotopological stacks are very nearly topological stacks:

Proposition 2.2. [19] *A stack $\mathcal{X}$ with representable diagonal is paratopological (pseudotopological) if and only if there exists a topological stack $\bar{\mathcal{X}}$ and a morphism $q : \bar{\mathcal{X}} \rightarrow \mathcal{X}$ such that for any paracompact space T (CW-complex T), q induces an equivalence of groupoids*

$$(1) \quad q(T) : \bar{\mathcal{X}}(T) \rightarrow \mathcal{X}(T).$$

The idea of the proof can be found in [19], but is enlightening, so we include it for completeness:

If q is as in (1), and $p : X \rightarrow \bar{\mathcal{X}}$ is an atlas for $\bar{\mathcal{X}}$, then $q \circ p : X \rightarrow \mathcal{X}$ satisfies condition ii) of Definition 2.6, hence $\mathcal{X}$ is paratopological. Conversely, if $\mathcal{X}$ is paratopological, take $p : X \rightarrow \mathcal{X}$ as in condition ii) of Definition 2.6. Form the weak fibred product

$$\begin{array}{ccc} X \times_{\mathcal{X}} X & \longrightarrow & X \\ \downarrow & & \downarrow p \\ X & \xrightarrow{p} & \mathcal{X}. \end{array}$$

Let $\bar{\mathcal{X}}$ be the topological stack associated with the topological groupoid

$$X \times_{\mathcal{X}} X \rightrightarrows X,$$

and $q : \bar{\mathcal{X}} \rightarrow \mathcal{X}$ the canonical map.

Definition 2.8. [19] A **homotopical** stack is a stack $\mathcal{X}$ on $\mathbf{TOP}$ (with respect to the open cover topology) with representable diagonal for which there exists a morphism $X \rightarrow \mathcal{X}$ from a topological space such that for all maps $T \rightarrow \mathcal{X}$ from T a CW-complex (equivalently T any space), the induced map $T \times_{\mathcal{X}} X \rightarrow T$ is a weak homotopy equivalence.

In [19], Noohi proved that paratopological and pseudotopological stacks are homotopical and that there is a functorial way of assigning them a map from a topological space satisfying property ii) of Definition 2.6 which is also a universal homotopy equivalence.

Proposition 2.3. [19] *Let $\mathcal{X}$ be a paratopological (pseudotopological) stack and $\bar{\mathcal{X}} \cong [\mathcal{G}]$ a topological stack as in Proposition 2.2. Then the composition*

$$\|\mathcal{G}\| \rightarrow \bar{\mathcal{X}} \rightarrow \mathcal{X}$$

is a universal weak equivalence. In particular, the weak homotopy type of $\mathcal{X}$ is the weak homotopy type of the topological stack $\bar{\mathcal{X}}$. The weak homotopy type of a homotopical stack is the weak homotopy type of the domain of the map $X \rightarrow \mathcal{X}$ defining it.

Definition 2.9. Any two stacks $\mathcal{X}$ and $\mathcal{Y}$ on $\mathbf{TOP}$ have an exponential stack $\mathcal{X}^{\mathcal{Y}}$ such that

$$\mathcal{X}^{\mathcal{Y}}(T) = \mathrm{Hom}(\mathcal{Y} \times T, \mathcal{X}).$$

We will from here on in denote $\mathcal{X}^{\mathcal{Y}}$ by $\mathrm{Map}(\mathcal{Y}, \mathcal{X})$ and refer to it as the **mapping stack** from $\mathcal{Y}$ to $\mathcal{X}$.

For the rest of this section, we work in the category $\mathbf{CG}$ of compactly generated spaces, which is Cartesian-closed.

In [20] Noohi proved:

Theorem 2.6. *If $\mathcal{X}$ and $\mathcal{Y}$ are topological stacks with $\mathcal{Y} \cong [\mathcal{H}]$ with $\mathcal{H}_0$ and $\mathcal{H}_1$ compact, then $\mathrm{Map}(\mathcal{Y}, \mathcal{X})$ is a topological stack.*

It appears that when $\mathcal{Y}$ does not satisfy this rather rigid compactness condition, that this may fail. However, Noohi also proved that when $\mathcal{Y}$ is instead locally compact, then $\mathrm{Map}(\mathcal{Y}, \mathcal{X})$ is at least paratopological:

Theorem 2.7. [20] *If $\mathcal{X}$ and $\mathcal{Y}$ are paratopological stacks with $\mathcal{Y} \cong [\mathcal{H}]$ such that $\mathcal{H}_0$ and $\mathcal{H}_1$ are locally compact, then $\mathrm{Map}(\mathcal{Y}, \mathcal{X})$ is a paratopological stack.*

We end this section by extending Noohi's result for the case when $\mathcal{Y} \cong [\mathcal{H}]$ and $\mathcal{H}_0$ is a CW-complex.

Remark. In what follows, we will further assume we are working with Hausdorff spaces.

Theorem 2.8. *If $\mathcal{X}$ and $\mathcal{Y}$ are topological stacks with $\mathcal{Y} \cong [\mathcal{H}]$ and $\mathcal{H}_0$ a CW-complex, then $\mathrm{Map}(\mathcal{Y}, \mathcal{X})$ is paratopological.*

Proof. Let $\mathcal{G}$ be a topological groupoid presenting $\mathcal{X}$. Let $\mathrm{Fib}(\mathcal{G})^{\mathcal{H}}$ denote the internal exponent of groupoid objects in compactly generated Hausdorff spaces. Let

$$\mathcal{K} := \left[\mathrm{Fib}(\mathcal{G})^{\mathcal{H}} \right].$$

Note that there is a canonical map

$$\varphi : \mathcal{K} \rightarrow \mathrm{Map}(\mathcal{Y}, \mathcal{X})$$

which sends any generalized homomorphism $T_{\mathcal{U}} \rightarrow \mathrm{Fib}(\mathcal{G})^{\mathcal{H}}$ to the induced generalized homomorphism from $T \times \mathcal{H}$, $T_{\mathcal{U}} \times \mathcal{H} \rightarrow \mathrm{Fib}(\mathcal{G})$ (which may be viewed as object in $\mathrm{Hom}_{\mathbf{Top}}(T \times \mathcal{Y}, \mathcal{X})$ since $\mathcal{G}$ and $\mathrm{Fib}(\mathcal{G})$ are Morita equivalent).

Suppose that T is a paracompact Hausdorff space. Then:

$$\begin{aligned}
\mathcal{K}(T) &\cong \varinjlim_{\mathcal{U}} \mathrm{Hom}_{\mathbb{C}\mathrm{GH}Gpd} \left(T_{\mathcal{U}}, \mathrm{Fib}(\mathcal{G})^{\mathcal{H}} \right) \\
&\cong \varinjlim_{\mathcal{U}} \mathrm{Hom}_{\mathbb{C}\mathrm{GH}Gpd} (T_{\mathcal{U}} \times \mathcal{H}, \mathrm{Fib}(\mathcal{G})).
\end{aligned}$$

Note that any paracompact Hausdorff space is a shrinking space so without loss of generality we may assume that each cover $\mathcal{U}$ of T is a topological covering by closed neighborhoods. Since any closed subset of a paracompact space is paracompact, this means that the groupoid $T_{\mathcal{U}}$ has paracompact object space. So the object space of $T_{\mathcal{U}} \times \mathcal{H}$ is the product of a paracompact Hausdorff space and a CW-complex, hence paracompact. Moreover, by Theorem 2.4, we have that

$$\mathrm{Hom}_{\mathbb{C}\mathrm{GH}Gpd} (T_{\mathcal{U}} \times \mathcal{H}, \mathrm{Fib}(\mathcal{G})) \cong \mathrm{Hom}_{\mathfrak{T}\mathfrak{S}\mathfrak{t}} (T \times \mathcal{Y}, \mathcal{X}).$$

Hence $\mathcal{K}(T) \cong \mathrm{Map}(\mathcal{Y}, \mathcal{X})(T)$. $\square$

3. THE COMPACTLY GENERATED GROTHENDIECK TOPOLOGY

3.1. The Compactly Generated Grothendieck Topology. Recall from section 1.1 that if Y is a compactly generated Hausdorff space and $\mathcal{X}$ a topological stack, then one might be tempted to claim that

$$\mathrm{Map}(Y, \mathcal{X}) := \varprojlim_{K_{\alpha} \hookrightarrow Y} \mathrm{Map}(K_{\alpha}, \mathcal{X}),$$

where the weak limit is taken over all compact subsets K_{α} of Y . However, there are some immediate problems with this temptation:

- This weak 2-limit may not exist as a topological stack.
- The fact that Y is the colimit of its compact subsets in $\mathbb{C}\mathrm{GH}$ does not imply that Y is the weak colimit of its compact subsets as a topological stack since the Yoneda embedding does not preserve arbitrary colimits.

Recall however that for an arbitrary subcanonical Grothendieck site $(\mathcal{C}, J)$, the Yoneda embedding $y : \mathcal{C} \hookrightarrow \mathrm{Sh}_J(\mathcal{C})$ preserves colimits of the form

$$C = \varinjlim_{C_{\alpha} \rightarrow C} C_{\alpha}$$

where $(C_{\alpha} \xrightarrow{f_{\alpha}} C)$ is a J -cover. We therefore shall construct a Grothendieck topology $\mathcal{CG}$ on $\mathbb{C}\mathrm{GH}$, called the compactly generated Grothendieck topology, such that for all Y , the inclusion of all compact subsets $(K_{\alpha} \hookrightarrow Y)$ is a $\mathcal{CG}$ -cover. As it shall turn out, in addition to being Cartesian closed, the 2-category of presentable stacks for this Grothendieck topology will also be complete.

In this subsection, we give a geometric construction of the compactly generated Grothendieck topology on $\mathbb{C}\mathrm{GH}$. Those readers not familiar with topos theory may wish to skip to Definition 3.1 for the concrete definition of a $\mathcal{CG}$ -cover. Some important properties of $\mathcal{CG}$ -covers are summarized as follows:

- Every open cover is a $\mathcal{CG}$ -cover (Proposition 3.2)
- For any space, the inclusion of all its compact subsets is a $\mathcal{CG}$ -cover (Corollary 3.1)

- iii) Every $\mathcal{CG}$ -cover of a locally compact space can be refined by an open one (Proposition 3.4)
- iv) The category of $\mathcal{CG}$ -sheaves over compactly generated Hausdorff spaces is equivalent to the category of ordinary sheaves over compact Hausdorff spaces (Theorem 3.2).

Let

$$j : \mathbb{CH} \hookrightarrow \mathbb{CGH}$$

be the full and faithful inclusion of compact Hausdorff spaces into compactly generated Hausdorff spaces. This induces a geometric morphism (j^*, j_*)

$$\mathrm{Set}^{\mathbb{CH}^{op}} \xrightleftharpoons[j_*]{j^*} \mathrm{Set}^{\mathbb{CGH}^{op}}$$

which is an embedding (i.e. j_* is full and faithful [10])².

Denote by

$$y_{\mathbb{CH}} : \mathbb{CGH} \rightarrow \mathrm{Set}^{\mathbb{CH}^{op}}$$

the functor which assigns a compactly generated Hausdorff space X the presheaf $T \mapsto \mathrm{Hom}_{\mathbb{CGH}}(T, X)$ and by

$$y_{\mathbb{CGH}} : \mathbb{CH} \rightarrow \mathrm{Set}^{\mathbb{CGH}^{op}}$$

the functor which assigns a $T \in \mathbb{CH}$ the presheaf $X \mapsto \mathrm{Hom}_{\mathbb{CGH}}(X, T)$. Note that $y_{\mathbb{CH}}$ is a fully faithful embedding. The pair (j^*, j_*) can be constructed as the adjoint pair induced by $y_{\mathbb{CH}}$ through a left Kan-extension \setminus nerve construction. Explicitly:

$$j^*(F)(T) = F(T)$$

and

$$j_*(G)(X) = \mathrm{Hom}_{\mathrm{Set}^{\mathbb{CH}^{op}}}(y_{\mathbb{CH}}(X), G).$$

From the general theory of adjoint functors, j_* restricts to an equivalence between, on one hand the full subcategory of $\mathrm{Set}^{\mathbb{CH}^{op}}$ whose objects are those for which the co-unit $\varepsilon(j)$ is an isomorphism, and on the other hand, the full subcategory of $\mathrm{Set}^{\mathbb{CGH}^{op}}$ whose objects are those for which the unit $\eta(j)$ is an isomorphism. However, since j is fully faithful, the co-unit is always an isomorphism, which verifies that j_* is fully faithful and gives us a way of describing its essential image.

In fact, j^* also has a left-adjoint $j_!$. The adjoint pair $j_! \dashv j^*$ is the one induced by $y_{\mathbb{CGH}}$. Hence, $j_!$ is the left Kan-extension of $y_{\mathbb{CGH}}$. We conclude that $j_!$ is also fully faithful (see: [10]).

Denote by $Sh(\mathbb{CGH})$ the topos of sheaves on compactly generated Hausdorff spaces with respect to the open cover topology. We define $Sh(\mathbb{CH})$ as the unique topos fitting in the following pullback diagram:

²Technically, these categories are not well defined due to set-theoretic issues, however, this can be overcome by careful use of Grothendieck universes. We will not dwell on this and all such similar size issues in this paper.

$$\begin{array}{ccc}
Sh(\mathbb{CH}) & \xrightarrow{\quad} & \text{Set}^{\mathbb{CH}^{op}} \\
\downarrow & & \downarrow \\
Sh(\mathbb{CGH}) & \xrightarrow{\quad} & \text{Set}^{\mathbb{CGH}^{op}}
\end{array}$$

Due to the factorization theorem of geometric morphisms in topos theory [10], the geometric embedding $Sh(\mathbb{CH}) \hookrightarrow \text{Set}^{\mathbb{CH}^{op}}$ corresponds to a Grothendieck topology $\mathcal{K}$ on $\mathbb{CH}$. It is easy to verify that since the functor j is fully faithful, the covering sieves in $\mathcal{K}$ for a compact Hausdorff space K are precisely those subobjects $S \rightrightarrows y(K)$ which are obtained by restricting a covering sieve of K with respect to the open cover topology on $\mathbb{CGH}$ to $\mathbb{CH}$ via the functor j^* . In this sense, the covering sieves are “the same as in the open cover topology”.

Proposition 3.1. *The Grothendieck topology $\mathcal{K}$ on $\mathbb{CH}$ has a basis of finite covers of the form $(T_i \hookrightarrow T)_{i=1}^n$ by compact neighborhoods (i.e. their interiors form an open cover).*

Proof. Compact Hausdorff spaces are locally compact in the strong sense that every point has a local base of compact neighborhoods. Hence covers by compact neighborhoods generate the same sieves as open covers. $\square$

Consider the geometric embedding

$$Sh(\mathbb{CH}) \xrightleftharpoons[i]{Sh} \text{Set}^{\mathbb{CH}^{op}},$$

where Sh denotes the sheafification with respect to $\mathcal{K}$.

Remark. It is clear that for any presheaf F in $\text{Set}^{\mathbb{CGH}^{op}}$, the $\mathcal{K}$ -sheafification of the restriction of F to $\mathbb{CH}$ is the same as the restriction of the sheafification of F .

By composition, we get an embedding of topoi

$$Sh(\mathbb{CH}) \xrightleftharpoons[j_* \circ i]{Sh \circ j^*} \text{Set}^{\mathbb{CGH}^{op}}.$$

Again by the factorization theorem [10], there exists a unique Grothendieck topology $\mathcal{CG}$ on $\mathbb{CGH}$ such that the category of sheaves $Sh_{\mathcal{CG}}(\mathbb{CGH})$ is $j_*(Sh(\mathbb{CH}))$. We will construct it and give some of its properties.

There is a very general construction [10] that shows how to extract the unique Grothendieck topology corresponding to this embedding.

First, we define a universal closure operation on $\text{Set}^{\mathbb{CGH}^{op}}$. Let F be a presheaf over $\mathbb{CGH}$ and let $m : A \rightrightarrows F$ be a representative for a subobject A of F . Then a representative for the subobject $\bar{A}$ is given by the left-hand side of the following pullback diagram

$$\begin{array}{ccc}
\bar{A} & \longrightarrow & j_* Sh j^*(A) \\
\downarrow & & \downarrow j_* Sh j^*(m) \\
F & \xrightarrow{\eta_F} & j_* Sh j^*(F),
\end{array}$$

where η is the unit of the adjunction $Sh \circ j^* \dashv j_* \circ i$.

To describe the covering sieves of $\mathcal{CG}$, it suffices to describe the universal closure operation on representables.

Let X be a compactly generated Hausdorff space.

Claim. The unit η_X is an isomorphism.

Proof. The restriction j^*X is a $\mathcal{K}$ -sheaf. Hence

$$j_*Shj^*X \cong j_*y_{\mathbb{CH}}(X).$$

Furthermore, for any compactly generated Hausdorff space Y ,

$$j_*y_{\mathbb{CH}}(X)(Y) = \text{Hom}_{\text{Set}^{\mathbb{CH}^{op}}}(y_{\mathbb{CH}}(Y), y_{\mathbb{CH}}(X)) \cong \text{Hom}_{\mathbb{CGH}}(Y, X)$$

since $y_{\mathbb{CH}}$ is fully faithful. $\square$

Now, let $m : A \rightarrow X$ a sieve. Then, since the unit η_X is an isomorphism, $\bar{A}$ is represented by the monomorphism

$$j_*Shj^*(A) \rightarrow X.$$

The covering sieves in $\mathcal{CG}$ of X are exactly those sieves on X whose closure is equal to the maximal sieve, i.e. X . So

$$m : A \rightarrow X$$

is a covering sieve if and only if

$$\bar{m} : j_*Shj^*(A) \rightarrow j_*j^*(X) \cong X$$

is an isomorphism. Since j_* is fully faithful, this is if and only if

$$\tilde{m} : Shj^*(A) \rightarrow j^*X$$

is an isomorphism. In other words,

$$m : A \rightarrow X$$

is a covering sieve if and only if the $\mathcal{K}$ -sheafification of $j^*(A)$ is isomorphic to $y_{\mathbb{CH}}(X)$

Definition 3.1. Let X be a compactly generated Hausdorff space and let

$$(\alpha_i : V_i \hookrightarrow X)_{i \in I}$$

be family of inclusions of subsets V_i of X . Such a family is called a $\mathcal{CG}$ -**cover** if for any compact subset K of X , there exists a (finite) subset $J(K) \subseteq I$ such that the collection $(V_j \cap K)_{j \in J(K)}$ can be refined by an open cover of K . Denote the set of $\mathcal{CG}$ -covers of X by $\mathcal{B}(X)$.

Lemma 3.1. $\mathcal{B}$ is a basis for the Grothendieck topology $\mathcal{CG}$.

Proof. Let $(f_i : Q_i \rightarrow X)$ be a class of maps into X . We denote the sieve it generates by S_f . For any compactly generated Hausdorff space Y , we have

$$S_f(Y) = \{h : Y \rightarrow X \text{ such that } h \text{ factors through } f_i \text{ for some } i\}.$$

So, S_f is a covering sieve if and only if when restricted to $\mathbb{CH}$, its $\mathcal{K}$ -sheafification is isomorphic to $y_{\mathbb{CH}}(X)$. We first note that $j^*(S_f)$ is clearly a $\mathcal{K}$ -separated presheaf. Hence, its sheafification is the same as $j^*(S_f)^+$. Since $y_{\mathbb{CH}}(X)$ is a sheaf, the canonical map

$$j^*(S_f) \rightarrow y_{\mathbb{CH}}(X)$$

factors uniquely as

$$\begin{array}{ccc} j^*(S_f) & \xrightarrow{\quad} & y_{\mathbb{CH}}(X) \\ \downarrow & \nearrow & \\ j^*(S_f)^+ & & \end{array}$$

It suffices to see when the map $j^*(S_f)^+ \rightarrow y_{\mathbb{CH}}(X)$ is an epimorphism. Let $\tilde{S}_f$ be the presheaf on $\mathbb{CH}$

$$\tilde{S}_f(T) = \left\{ \mathcal{U} = (U_i)_{i=1}^n, (a_j \in S_f(U_j))_{j=1}^n \mid a_i|_{U_{ij}} = a_j|_{U_{ij}} \text{ for all } i, j \right\}.$$

Then the map $j^*(S_f)^+ \rightarrow y_{\mathbb{CH}}(X)$ fits in a diagram

$$\begin{array}{ccc} \tilde{S}_f & \xrightarrow{\quad} & y_{\mathbb{CH}}(X) \\ \downarrow & \nearrow & \\ j^*(S_f)^+ & & \end{array}$$

It suffices to see when the canonical map $\tilde{S}_f \rightarrow y_{\mathbb{CH}}(X)$ is pointwise surjective. This is precisely when for any map $h : K \rightarrow X$ from a compact space $K \in \mathbb{CH}$, there exists an open cover $(U_j)_j$ of K such that for all j , $h|_{U_j}$ factors through f_i for some i . Classes of maps with codomain X with this property constitute a large basis for the Grothendieck topology $\mathcal{CG}$. It is in fact maximal in the sense that S is a covering sieve if and only if it is one generated by a large cover of this form. We will now show that any such large covering family has a refinement by one of the form of the lemma.

Let $(f_i : Q_i \rightarrow X)$ denote such a (possibly large) family and let

$$(i_\alpha : K_\alpha \hookrightarrow X)$$

denote the inclusion of all compact subsets of X . Then for each α , there exists a finite open cover of K_α , $(\mathcal{O}_j^\alpha)$ such that the inclusion of each $\mathcal{O}_j^\alpha$ into X factors through some

$$f_{j,\alpha} : Q_{j,\alpha} \rightarrow X$$

via a map

$$\lambda_j^\alpha : \mathcal{O}_j^\alpha \rightarrow Q_{j,\alpha}.$$

Let $\mathcal{U} := (\mathcal{O}_j^\alpha \hookrightarrow X)_{j,\alpha}$. Let $g : L \rightarrow X$ be a map with $L \in \mathbb{CH}$. Then $g(L) = K_\alpha$ for some α . Let

$$\mathcal{V}_L^g := (g^{-1}(\mathcal{O}_j^\alpha)).$$

Then $\mathcal{V}_L^g$ is an open cover of L such that the restriction of g to any element of the cover factors through the inclusion of some $\mathcal{O}_j^\alpha$ into X . Hence the sieve generated by $\mathcal{U}$ is a covering sieve for $\mathcal{CG}$ which refines the sieve generated by $(f_i : Q_i \rightarrow X)$. $\square$

We have the following obvious proposition whose converse is not true:

Proposition 3.2. *Any open cover of a space is also a $\mathcal{CG}$ cover.*

In particular, one cover that is quite useful is the following.

Corollary 3.1. *For any compactly generated Hausdorff space, the inclusion of all compact subsets is a $\mathcal{CG}$ -cover.*

However, for the category $\mathbb{LCH}$ of locally compact Hausdorff spaces, it suffices to work with open covers:

Proposition 3.3. *Every $\mathcal{CG}$ -cover of a locally compact Hausdorff space $X \in \mathbb{LCH}$ can be refined by an open covering.*

Proof. Let $X \in \mathbb{LCH}$ and let $\mathcal{V} = (\alpha_i : V_i \hookrightarrow X)_{i \in I}$ be a $\mathcal{CG}$ -cover of X . Let (K_l) be a topological covering of X by compact subsets such that the interiors $\text{int}(K_l)$ constitute an open cover for X . Then for each K_l , there exists a finite subset $J(K_l) \subseteq I$ such that

$$(V_j \cap K_l)_{j \in J(K_l)}$$

is an open cover for K_l such that the inclusion of each $V_j \cap K_l$ into X factors through the inclusion of V_j . Let $\mathcal{U} := (V_j \cap \text{int}(K_l))_{l, j \in J(K_l)}$. Then $\mathcal{U}$ is an open cover of X refining $\mathcal{V}$. $\square$

We can now define the $\mathcal{CG}$ -sheafification functor $Sh_{\mathcal{CG}}$ either by the covering sieves, or by using the basis $\mathcal{B}$ (i.e. both will give naturally isomorphic functors). Let $Sh_{\mathcal{CG}}(\mathbb{CGH})$ denote the category of $\mathcal{CG}$ -sheaves. Then we have an embedding of topoi given by

$$Sh_{\mathcal{CG}}(\mathbb{CGH}) \xrightleftharpoons[\ell]{Sh_{\mathcal{CG}}} \text{Set}^{\mathbb{CGH}^{op}},$$

where $\ell : Sh_{\mathcal{CG}}(\mathbb{CGH}) \hookrightarrow \text{Set}^{\mathbb{CGH}^{op}}$ is the inclusion of the category of sheaves.

By the previous observation that open covers are $\mathcal{CG}$ -covers, we also have

$$Sh_{\mathcal{CG}}(\mathbb{CGH}) \subset Sh(\mathbb{CGH}),$$

where $Sh(\mathbb{CGH})$ is the category of sheaves on $\mathbb{CGH}$ with respect to the open cover topology.

By construction, we have the following theorem:

Theorem 3.2. *There is an equivalence of topoi*

$$Sh(\mathbb{CH}) \xrightleftharpoons[Sh_{\mathcal{CG}} \circ j_* \circ i]{Sh \circ j^* \circ \ell} Sh_{\mathcal{CG}}(\mathbb{CGH})$$

such that

$$Sh(\mathbb{CH}) \xrightleftharpoons[Sh_{\mathcal{CG}} \circ j_* \circ i]{Sh \circ j^* \circ \ell} Sh_{\mathcal{CG}}(\mathbb{CGH}) \xrightleftharpoons[\ell]{Sh_{\mathcal{CG}}} \text{Set}^{\mathbb{CGH}^{op}}$$

is a factorization of

$$Sh(\mathbb{CH}) \xrightleftharpoons[j_* \circ i]{Sh \circ j^*} \text{Set}^{\mathbb{CGH}^{op}}$$

(up to natural isomorphism).

Note that the essential image of ℓ is the same as the essential image of $j_* \circ i$. Hence, a presheaf F in $\text{Set}^{\mathbb{C}\mathbb{G}\mathbb{H}^{op}}$ is a $\mathcal{CG}$ -sheaf if and only if the unit η of $Sh \circ j^* \dashv j_* \circ i$ is an isomorphism at F . We have the following immediate corollary.

Corollary 3.2. *The Grothendieck topology $\mathcal{CG}$ is subcanonical.*

Lemma 3.3. *If F is a $\mathcal{CG}$ -sheaf, then $j^*(F)$ is a $\mathcal{X}$ -sheaf.*

Proof. Since F is a $\mathcal{CG}$ -sheaf, it is in the essential image of $j_* \circ i$, hence, via η_F , we have

$$F \cong j_* Sh j^*(F).$$

By applying j^* we have

$$j^*(F) \cong j^* j_* Sh j^*(F).$$

Since the co-unit $\varepsilon(j)$ of $j^* \dashv j_*$ is an isomorphism, this yields

$$j^*(F) \cong Sh j^*(F).$$

□

Corollary 3.3. *If F is a presheaf in $\text{Set}^{\mathbb{C}\mathbb{G}\mathbb{H}^{op}}$,*

$$Sh_{\mathcal{CG}}(F) \cong j_* Sh(j^*F).$$

If $F \in Sh(\mathbb{C}\mathbb{G}\mathbb{H})$ is a sheaf in the open cover topology then its $\mathcal{CG}$ -sheafification is given by $j_ j^* F$.*

Proof.

$$j_* Sh(j^*F) \cong (j_* \circ i) \circ Sh(j^*F),$$

so $j_* Sh(j^*F)$ is in the image of $j_* \circ i$, hence a $\mathcal{CG}$ -sheaf. Now, let G be any $\mathcal{CG}$ -sheaf. Then:

$$\begin{aligned} \text{Hom}_{\text{Set}^{\mathbb{C}\mathbb{G}\mathbb{H}^{op}}}(j_* Sh(j^*F), G) &\cong \text{Hom}_{\text{Set}^{\mathbb{C}\mathbb{H}^{op}}}(Sh j^*(F), j^*(G)) \\ &\cong \text{Hom}_{\text{Set}^{\mathbb{C}\mathbb{H}^{op}}}(j^*(F), j^*(G)) \\ &\cong \text{Hom}_{\text{Set}^{\mathbb{C}\mathbb{G}\mathbb{H}^{op}}}(F, j_* j^*(G)) \\ &\cong \text{Hom}_{\text{Set}^{\mathbb{C}\mathbb{G}\mathbb{H}^{op}}}(F, G). \end{aligned}$$

□

We end this subsection by noting ordinary sheaves and $\mathcal{CG}$ -sheaves agree on locally compact Hausdorff spaces:

Let ς denote the unit of the adjunction $j^* \dashv j_*$.

Proposition 3.4. *Let $F \in Sh(\mathbb{C}\mathbb{G}\mathbb{H})$ be a sheaf in the open cover topology and X in $\mathbb{LCH}$ a locally compact Hausdorff space. Then the map*

$$\varsigma_F(X) : F(X) \rightarrow j_* j^* F(X) \cong Sh_{\mathcal{CG}}(F)(X)$$

is a bijection. In particular, F and $Sh_{\mathcal{CG}}(F)$ agree on locally compact Hausdorff spaces.

Proof. This follows immediately from Proposition 3.3

□

3.2. Stacks for the Compactly Generated Grothendieck Topology. Denote again by $y_{\mathbb{CH}}$ the 2-functor

$$y_{\mathbb{CH}} : \mathbb{CGH} \rightarrow Gpd^{\mathbb{CH}^{op}}$$

induced by the inclusion

$$(\cdot)^{(id)} : \text{Set}^{\mathbb{CH}^{op}} \hookrightarrow Gpd^{\mathbb{CH}^{op}}.$$

Then, it produces a 2-adjoint pair, which we will also denote by $j^* \dashv j_*$, by constructing j^* as the weak left Kan extension of $y_{\mathbb{CH}}$, and by letting

$$j_* \mathcal{Y}(X) := \text{Hom}_{Gpd^{\mathbb{CH}^{op}}}(y_{\mathbb{CH}}(X), \mathcal{Y}).$$

By setting

$$j^*(\mathcal{X})(T) := \mathcal{X}(T)$$

we get a 2-functor which is weak-colimit preserving and whose restriction to representables is the same as $y_{\mathbb{CH}}$, hence, by uniqueness, the above equation for j^* must be correct. Note that the co-unit is an equivalence, hence j_* is fully faithful. Similarly, denote again by $y_{\mathbb{CGH}}$ the 2-functor

$$y_{\mathbb{CGH}} : \mathbb{CH} \rightarrow Gpd^{\mathbb{CGH}^{op}}$$

induced by the inclusion

$$(\cdot)^{(id)} : \text{Set}^{\mathbb{CGH}^{op}} \hookrightarrow Gpd^{\mathbb{CGH}^{op}}.$$

Let $j_!$ be the 2-functor obtained by its weak left Kan extension. Just as before, $j_!$ is left 2-adjoint to j^* . Similarly, $j_!$ is fully faithful.

To justify the use of the same symbols, we have

$$\begin{aligned} (\cdot)^{(id)} \circ j^* &= j^* \circ (\cdot)^{(id)} \\ (\cdot)^{(id)} \circ j_* &= j_* \circ (\cdot)^{(id)} \\ (\cdot)^{(id)} \circ j_! &= j_! \circ (\cdot)^{(id)} \end{aligned}$$

where, the j^* , j_* , $j_!$ appearing on the left-hand side are 1-functors.

Let $St(\mathbb{CH})$ denote the 2-category of stacks on $\mathbb{CH}$ with respect to the Grothendieck topology $\mathcal{K}$. Then we have a 2-adjoint pair $St \dashv i$

$$St(\mathbb{CH}) \xrightleftharpoons[i]{St} Gpd^{\mathbb{CH}^{op}},$$

where St is the stackification 2-functor (Definition A.3) and i is the inclusion. Then, by composition, we get a 2-adjoint pair $St \circ j^* \dashv j_* \circ i$

$$St(\mathbb{CH}) \xrightleftharpoons[j_* \circ i]{St \circ j^*} Gpd^{\mathbb{CGH}^{op}}.$$

Definition 3.2. A stack with respect to the Grothendieck topology $\mathcal{CG}$ on $\mathbb{CGH}$ will be called a $\mathcal{CG}$ -stack.

Let $St_{\mathcal{CG}}(\mathbb{CGH})$ denote full sub-2-category of $Gpd^{\mathbb{CGH}^{op}}$ consisting of $\mathcal{CG}$ -stacks, and let $St_{\mathcal{CG}}$ denote the associated stackification 2-functor, and ℓ the inclusion, so $St_{\mathcal{CG}} \dashv \ell$.

Just as before, since every open covering is a $\mathcal{CG}$ -cover,

$$St_{\mathcal{CG}}(\mathbb{CGH}) \subset St(\mathbb{CGH}).$$

The following results and their proofs follow naturally from those of the previous section when combined with the Comparison Lemma for stacks, a straight-forward stacky analogue of the theorem in [23] III:

Corollary 3.4. *There is an equivalence of 2-categories*

$$St(\mathbb{CH}) \xrightleftharpoons[St_{\mathcal{CG}} \circ j_* \circ i]{St \circ j^* \circ \ell} St_{\mathcal{CG}}(\mathbb{CGH}),$$

such that

$$St(\mathbb{CH}) \xrightleftharpoons[St_{\mathcal{CG}} \circ j_* \circ i]{St \circ j^* \circ \ell} St_{\mathcal{CG}}(\mathbb{CGH}) \xrightleftharpoons[\ell]{St_{\mathcal{CG}}} Gpd^{\mathbb{CGH}^{op}}$$

is a factorization of

$$St(\mathbb{CH}) \xrightleftharpoons[j_* \circ i]{St \circ j^*} Gpd^{\mathbb{CGH}^{op}}$$

(up to natural equivalence).

Lemma 3.4. *If $\mathcal{X}$ is a $\mathcal{CG}$ -stack, then $j^*(\mathcal{X})$ is a $\mathcal{K}$ -stack.*

Corollary 3.5. *If $\mathcal{X}$ is a prestack in $Gpd^{\mathbb{CGH}^{op}}$ (see Appendix A for terminology),*

$$St_{\mathcal{CG}}(\mathcal{X}) \cong j_* St(j^* \mathcal{X}).$$

If $\mathcal{X} \in St(\mathbb{CGH})$ is a stack in the open cover topology then its $\mathcal{CG}$ -stackification is given by $j_ j^* \mathcal{X}$.*

Again, let ς denote the unit of the 2-adjunction $j^* \dashv j_*$.

Proposition 3.5. *Let $\mathcal{X} \in St(\mathbb{CGH})$ be a stack in the open cover topology and X in $\mathbb{LCH}$ a locally compact Hausdorff space. Then the map*

$$\varsigma_{\mathcal{X}}(X) : \mathcal{X}(X) \rightarrow j_* j^* \mathcal{X}(X) \cong St_{\mathcal{CG}}(\mathcal{X})(X)$$

is an equivalence of groupoids. In particular, $\mathcal{X}$ and $St_{\mathcal{CG}}(\mathcal{X})$ agree on locally compact Hausdorff spaces.

4. COMPACTLY GENERATED STACKS

4.1. Compactly Generated Stacks.

Definition 4.1. A **compactly generated stack** is a presentable $\mathcal{CG}$ -stack (see Definition B.16).

We denote the full sub-2-category of $St_{\mathcal{CG}}(\mathbb{CGH})$ of compactly generated stacks by $\mathcal{CG}\mathfrak{IS}\mathfrak{t}$.

Intrinsically, a compactly generated stack is a $\mathcal{CG}$ -stack $\mathcal{X}$ such that there exists a compactly generated Hausdorff space X and a representable $\mathcal{CG}$ -epimorphism

$$p : X \rightarrow \mathcal{X}.$$

The map above is a $\mathcal{CG}$ -**atlas** for $\mathcal{X}$.

Let

$$\hat{y}_{\mathbb{CH}} : \mathbb{CH}Gpd \rightarrow Gpd^{\mathbb{CH}^{op}}$$

denote the 2-functor

$$\mathcal{G} \mapsto \text{Hom}_{\mathbb{C}\mathcal{G}\mathcal{H}Gpd} \left((\cdot)^{id}, \mathcal{G} \right).$$

Given a topological groupoid $\mathcal{G}$ in $\mathbb{C}\mathcal{G}\mathcal{H}Gpd$, denote by $[\mathcal{G}]_{\mathcal{K}}$ the associated $\mathcal{K}$ -stack $St \circ \tilde{y}_{\mathbb{C}\mathcal{H}}(\mathcal{G})$.

Let $\mathcal{C}\mathcal{G}\mathcal{I}\mathcal{S}\mathfrak{t}'$ denote the essential image in $Gpd^{\mathbb{C}\mathcal{H}op}$ of $St \circ \tilde{y}_{\mathbb{C}\mathcal{H}}$, i.e., it is the full sub-2-category of consisting of $\mathcal{K}$ -stacks equivalent to $[\mathcal{G}]_{\mathcal{K}}$ for some compactly generated topological groupoid $\mathcal{G}$. It is immediate from Theorem 3.4 that this 2-category is equivalent to $\mathcal{C}\mathcal{G}\mathcal{I}\mathcal{S}\mathfrak{t}$. In fact, the functor j_* restricts to an equivalence $j_* : \mathcal{C}\mathcal{G}\mathcal{I}\mathcal{S}\mathfrak{t}' \rightarrow \mathcal{C}\mathcal{G}\mathcal{I}\mathcal{S}\mathfrak{t}$ of 2-categories. Hence we have proven:

Theorem 4.1. *The 2-category of compactly generated stacks, $\mathcal{C}\mathcal{G}\mathcal{I}\mathcal{S}\mathfrak{t}$, is equivalent to the essential image of*

$$j^* : \mathcal{I}\mathcal{S}\mathfrak{t} \rightarrow St(\mathbb{C}\mathcal{H}).$$

Note that from Theorems B.1 and B.2, $\mathcal{C}\mathcal{G}\mathcal{I}\mathcal{S}\mathfrak{t}$ is also equivalent to the bicategory of fractions $\mathbb{C}\mathcal{G}\mathcal{H}Gpd[W_{\mathcal{C}\mathcal{G}}^{-1}]$ of compactly generated Hausdorff topological groupoids with inverted $\mathcal{C}\mathcal{G}$ -Morita equivalences, and also to the bicategory $Bun^{\mathcal{C}\mathcal{G}}\mathbb{C}\mathcal{G}\mathcal{H}Gpd$ of compactly generated Hausdorff topological groupoids with left $\mathcal{C}\mathcal{G}$ -principal bundles as morphisms:

Theorem 4.2. *The 2-functor*

$$St_{\mathcal{C}\mathcal{G}} \circ \tilde{y} : \mathbb{C}\mathcal{G}\mathcal{H}Gpd \rightarrow \mathcal{C}\mathcal{G}\mathcal{I}\mathcal{S}\mathfrak{t}$$

induces an equivalence of bicategories

$$P_{\mathcal{C}\mathcal{G}} : \mathbb{C}\mathcal{G}\mathcal{H}Gpd[W_{\mathcal{C}\mathcal{G}}^{-1}] \xrightarrow{\sim} \mathcal{C}\mathcal{G}\mathcal{I}\mathcal{S}\mathfrak{t}.$$

Theorem 4.3. *The 2-functor*

$$St_{\mathcal{C}\mathcal{G}} \circ \tilde{y} : \mathbb{C}\mathcal{G}\mathcal{H}Gpd \rightarrow \mathcal{C}\mathcal{G}\mathcal{I}\mathcal{S}\mathfrak{t}$$

induces an equivalence of bicategories

$$P'_{\mathcal{C}\mathcal{G}} : Bun^{\mathcal{C}\mathcal{G}}\mathbb{C}\mathcal{G}\mathcal{H}Gpd \xrightarrow{\sim} \mathcal{C}\mathcal{G}\mathcal{I}\mathcal{S}\mathfrak{t}.$$

We note that the principal bundles in $Bun^{\mathcal{C}\mathcal{G}}\mathbb{C}\mathcal{G}\mathcal{H}Gpd$ have a very simple description:

Recall that our notion of principal bundle depends on a Grothendieck topology. When the projection map of a principal bundle admits local sections (with respect to the open cover topology), it is called **ordinary**.

Proposition 4.1. *If $\mathcal{G}$ is a topological groupoid in $\mathbb{C}\mathcal{G}\mathcal{H}Gpd$, X is a compactly generated Hausdorff space, and*

$$\begin{array}{ccc} \mathcal{G}_1 & \xrightarrow{\mathcal{C}} & P \\ \downarrow & \mu \swarrow & \downarrow \pi \\ \mathcal{G}_0 & & X \end{array}$$

is a left $\mathcal{G}$ -space over π , then it is a $\mathcal{C}\mathcal{G}$ -principal bundle if and only if the restriction of P to K is an ordinary principal $\mathcal{G}$ -bundle over K , for every compact subset $K \subseteq X$.

Proof. Suppose that we are given a left $\mathcal{G}$ -space over π, P , whose restriction to K is an ordinary principal $\mathcal{G}$ -bundle over K , for every compact subset $K \subseteq X$. Then, for each compact subset $K_\alpha \subseteq X$, we can choose an open cover $(U_\alpha^i \hookrightarrow K_\alpha)_{i=1}^{N_\alpha}$ over which P admits local sections. Then P admits local sections with respect to the $\mathcal{G}$ -cover

$$\mathcal{U} := (U_\alpha^i \hookrightarrow X).$$

The converse is trivial. $\square$

Corollary 4.1. *If $\mathcal{G}$ and $\mathcal{H}$ are topological groupoids in $\mathbb{C}\mathbb{G}\mathbb{H}\mathbb{G}pd$ and $\mathcal{H}_0$ is locally compact, then*

$$Bun_{\mathcal{G}}^{\mathcal{G}}(\mathcal{H}) \cong Bun_{\mathcal{G}}(\mathcal{H}),$$

where $Bun_{\mathcal{G}}^{\mathcal{G}}(\mathcal{H})$ denotes the groupoid of $\mathcal{G}$ -principal $\mathcal{G}$ -bundles over $\mathcal{H}$, and $Bun_{\mathcal{G}}(\mathcal{H})$ denotes the groupoid of ordinary principal $\mathcal{G}$ -bundles over $\mathcal{H}$.

Equivalently:

If $\mathcal{X}$ and $\mathcal{Y}$ are topological stacks, and $\mathcal{Y}$ admits a locally compact Hausdorff atlas $Y \rightarrow \mathcal{Y}$, then the map

$$\mathrm{Hom}_{\mathrm{St}(\mathbb{C}\mathbb{G}\mathbb{H})}(\mathcal{Y}, \mathcal{X}) \rightarrow \mathrm{Hom}_{\mathrm{St}(\mathbb{C}\mathbb{G}\mathbb{H})}(St_{\mathcal{G}}(\mathcal{Y}), St_{\mathcal{G}}(\mathcal{X}))$$

induced by the unit $\varsigma_{\mathcal{X}} : \mathcal{X} \rightarrow St_{\mathcal{G}}(\mathcal{X})$, and the 2-adjunction $St_{\mathcal{G}} \dashv \ell$ is an equivalence of groupoids.

Theorem 4.4. *The 2-category of compactly generated stacks is equivalent to the sub-2-category of topological stacks consisting of those topological stacks which admit a locally compact atlas.³*

Proof. First note that the sub-2-category of topological stacks consisting of those topological stacks which admit a locally compact atlas is equivalent to the sub-bicategory of topological groupoids and ordinary principal bundles consisting of topological groupoids with locally compact object space. Denote this bicategory by $Bun\mathbb{L}\mathbb{C}\mathbb{H}_0\mathbb{G}pd$. Consider now the map of bicategories

$$\Upsilon : Bun\mathbb{L}\mathbb{C}\mathbb{H}_0\mathbb{G}pd \rightarrow \mathcal{G}\mathcal{G}\mathbb{T}\mathbb{S}\mathfrak{t}$$

which sends the groupoid $\mathcal{G}$ to $[\mathcal{G}]_{\mathcal{G}}$. Note that if $\mathcal{G}$ and $\mathcal{H}$ are in $Bun\mathbb{L}\mathbb{C}\mathbb{H}_0\mathbb{G}pd$,

$$\begin{aligned} \mathrm{Hom}_{Bun\mathbb{L}\mathbb{C}\mathbb{H}_0\mathbb{G}pd}(\mathcal{H}, \mathcal{G}) &= Bun_{\mathcal{G}}(\mathcal{H}) \\ &\cong Bun_{\mathcal{G}}^{\mathcal{G}}(\mathcal{H}) \\ &\cong \mathrm{Hom}_{\mathcal{G}\mathcal{G}\mathbb{T}\mathbb{S}\mathfrak{t}}([\mathcal{H}]_{\mathcal{G}}, [\mathcal{G}]_{\mathcal{G}}) \\ &= \mathrm{Hom}_{\mathcal{G}\mathcal{G}\mathbb{T}\mathbb{S}\mathfrak{t}}(\Upsilon(\mathcal{H}), \Upsilon(\mathcal{G})), \end{aligned}$$

since $\mathcal{H}_0$ is locally compact. Hence Υ is fully faithful. Furthermore, notice the essential image of Υ is those compactly generated stacks which admit a locally compact atlas. To complete the proof, note that if $\mathcal{G}$ is any topological groupoid, then the inclusion of all compact subsets of $\mathcal{G}_0$ is a $\mathcal{G}$ -cover $\mathcal{U}$ and hence $\mathcal{G}_{\mathcal{U}}$ is a topological groupoid with locally compact object space which is $\mathcal{G}$ -Morita

³When we work over compactly generated Hausdorff spaces

equivalent to $\mathcal{G}$. Hence, every compactly generated stack admits a locally compact atlas and $\mathfrak{Y}$ is essentially surjective. $\square$

Theorem 4.5. *The 2-category of compactly generated stacks has arbitrary products.*

Proof. Let $\mathcal{X}_i$ be an arbitrary family of compactly generated stacks. Then we can choose topological groupoids $\mathcal{G}_i$ in $\mathbb{C}\mathbb{G}\mathbb{H}Gpd$ such that

$$\mathcal{X}_i \cong [\mathcal{G}_i]_{\mathcal{C}\mathcal{G}}.$$

Note that

$$[\mathcal{G}_i]_{\mathcal{C}\mathcal{G}} \cong j_* [\mathcal{G}_i]_{\mathcal{K}}.$$

In light of Lemma 2.3, we may assume without loss of generality that each $\mathcal{G}_i$ is fibrant. Under this assumption, by Theorem 2.4, it follows that

$$[\mathcal{G}_i]_{\mathcal{C}\mathcal{G}} \cong j_* \tilde{y}_{\mathbb{C}\mathbb{H}}(\mathcal{G}_i).$$

Note that the product $\prod_i \mathcal{X}_i$ is a $\mathcal{C}\mathcal{G}$ -stack, as any 2-category of stacks is complete. It suffices to show that this product is still presentable.

Recall that $\tilde{y}_{\mathbb{C}\mathbb{H}}$ preserves small weak limits. Moreover, j_* does as well as it is a right 2-adjoint. Hence

$$\prod_i \mathcal{X}_i \cong \prod_i j_* \tilde{y}_{\mathbb{C}\mathbb{H}}(\mathcal{G}_i) \cong j_* \tilde{y}_{\mathbb{C}\mathbb{H}}\left(\prod_i \mathcal{G}_i\right).$$

It follows that

$$\begin{aligned} \prod_i \mathcal{X}_i &\cong St_{\mathcal{C}\mathcal{G}}\left(\prod_i \mathcal{X}_i\right) \cong St_{\mathcal{C}\mathcal{G}}\left(j_* \tilde{y}_{\mathbb{C}\mathbb{H}}\left(\prod_i \mathcal{G}_i\right)\right) \\ &\cong (j_* \circ St \circ j^*) \circ j_* \circ (j^* \tilde{y})\left(\prod_i \mathcal{G}_i\right) \\ &\cong (j_* \circ St \circ j^*) \circ (\tilde{y})\left(\prod_i \mathcal{G}_i\right) \\ &\cong \left[\prod_i \mathcal{G}_i\right]_{\mathcal{C}\mathcal{G}}. \end{aligned}$$

$\square$

Corollary 4.2. *The 2-category of compactly generated stacks is closed under arbitrary small weak limits.*

Proof. Since $\mathbb{C}\mathbb{G}\mathbb{H}Gpd$ is closed under binary weak fibred products and the stackification 2-functor $St_{\mathcal{C}\mathcal{G}}$ preserves finite weak limits, the 2-category $\mathcal{C}\mathcal{G}\mathfrak{I}\mathfrak{S}\mathfrak{t}$ is closed under binary weak fibred products. By Theorem 4.5, this 2-category has arbitrary small products. Hence, again by [25] and [6], $\mathcal{C}\mathcal{G}\mathfrak{I}\mathfrak{S}\mathfrak{t}$ is a complete 2-category. $\square$

4.2. Mapping Stacks of Compactly Generated Stacks. Recall that if $\mathcal{X}$ and $\mathcal{Y}$ are any stacks over $\mathbb{C}\mathbb{H}$ then they have a mapping stack

$$\mathrm{Map}(\mathcal{Y}, \mathcal{X})(T) = \mathrm{Hom}_{Gpd^{\mathbb{C}\mathbb{H}^{op}}}(\mathcal{Y} \times T, \mathcal{X}).$$

It is the goal of this section to prove that if $\mathcal{X}$ and $\mathcal{Y}$ are compactly generated stacks, then so is $\mathrm{Map}(\mathcal{Y}, \mathcal{X})$.

Lemma 4.6. *If $\mathcal{Y} \cong [\mathcal{H}]_{\mathcal{CG}}$ is a compactly generated stack with $\mathcal{H}_0$ and $\mathcal{H}_1$ compact, and $\mathcal{X} \cong [\mathcal{G}]_{\mathcal{CG}}$ an arbitrary compactly generated stack, then $\mathrm{Map}(\mathcal{Y}, \mathcal{X})$ is a compactly generated stack. More specifically, if $\mathcal{K}$ is a presentation for the topological stack $\mathrm{Map}([\mathcal{H}], [\mathcal{G}])$ ensured by Theorem 2.6, then*

$$\mathrm{Map}(\mathcal{Y}, \mathcal{X}) \cong [\mathcal{K}]_{\mathcal{CG}}.$$

In particular, $\mathrm{Map}(\mathcal{Y}, \mathcal{X})$ and $\mathrm{Map}([\mathcal{H}], [\mathcal{G}])$ restrict to the same stack over locally compact Hausdorff spaces.

Proof. Since any $\mathcal{CG}$ -stack is completely determined by its restriction to $\mathbb{C}\mathbb{H}$, it suffices to show that for any compact Hausdorff space $T \in \mathbb{C}\mathbb{H}$,

$$[\mathcal{K}]_{\mathcal{CG}}(T) \cong \mathrm{Hom}_{Gpd^{\mathbb{C}\mathbb{H}^{op}}}(\mathcal{Y} \times T, \mathcal{X}).$$

But, since T is compact Hausdorff

$$[\mathcal{K}]_{\mathcal{CG}}(T) \cong [\mathcal{K}](T)$$

and because of the definition of $\mathcal{K}$

$$[\mathcal{K}](T) \cong \mathrm{Hom}_{Gpd^{\mathbb{C}\mathbb{H}^{op}}}(\mathcal{Y} \times T, [\mathcal{G}]).$$

Furthermore, since $\mathcal{Y} \times T \cong [\mathcal{H} \times T]$ and $\mathcal{H} \times T$ has compact Hausdorff object space, by Corollary 4.1,

$$\mathrm{Hom}_{Gpd^{\mathbb{C}\mathbb{H}^{op}}}(\mathcal{Y} \times T, [\mathcal{G}]) \cong \mathrm{Hom}_{Gpd^{\mathbb{C}\mathbb{H}^{op}}}(\mathcal{Y} \times T, \mathcal{X}).$$

□

Corollary 4.3. *If K is a compact Hausdorff space and $\mathcal{X}$ an arbitrary compactly generated stack, then $\mathrm{Map}(K, \mathcal{X})$ is a compactly generated stack.*

Lemma 4.7. *If X is a compactly generated Hausdorff space and $\mathcal{X}$ an arbitrary compactly generated stack, then $\mathrm{Map}(X, \mathcal{X})$ is a compactly generated stack.*

Proof. Let $(K_\alpha \xrightarrow{i_\alpha} X)$ denote the inclusion of all compact subsets of X . This is a $\mathcal{CG}$ -cover for X . Let Y be an arbitrary compactly generated Hausdorff space. Note that $(K_\alpha \times Y \xrightarrow{i_\alpha \times id_Y} X \times Y)$ is also a $\mathcal{CG}$ -cover.

By Proposition A.1, we have that in $\mathcal{CG}\mathfrak{T}\mathfrak{S}\mathfrak{t}$

$$X \cong \varinjlim_{K_\alpha \hookrightarrow X} K_\alpha$$

and

$$X \times Y \cong \varinjlim_{K_\alpha \times Y \hookrightarrow X \times Y} (K_\alpha \times Y).$$

Hence

Remark. This implies that the 2-category of topological stacks (in compactly generated Hausdorff spaces) with locally compact atlases is Cartesian-closed. This might seem surprising since, after all, locally compact Hausdorff spaces are quite far from

being Cartesian closed. What is happening is that the exponential of two locally compact spaces is not a space in this description, but a stack! In fact, the category of compactly generated Hausdorff spaces embeds into this 2-category by sending a space X to the stack associated to the topological groupoid $(X)_{\mathcal{K}}^{(id)}$, where $\mathcal{K}$ denotes the $\mathcal{CG}$ -cover which is the inclusion of all compact subsets of X .

4.3. Homotopy Type of Compactly Generated Stacks. In this section, we show that every compactly generated stacks admits a $\mathcal{CG}$ -atlas which is a universal weak equivalence. As a Corollary, we show that the homotopy type of the classifying space of a topological groupoid is invariant under $\mathcal{CG}$ -Morita equivalences.

Remark. The notion of universal weak equivalence is local on the target with respect to the Grothendieck topology $\mathcal{CG}$. This is a consequence of the fact that the singular nerve of a topological space is the colimit of the singular nerve of each of its compact subsets.

Theorem 4.9. *Let $\mathcal{X} \cong [\mathcal{G}]_{\mathcal{CG}}$ be a compactly generated stack. Then the atlas $\|\mathcal{G}\| \rightarrow \mathcal{X}$ is a universal weak equivalence.*

Proof. In [19], it is shown that we have a 2-Cartesian diagram

$$\begin{array}{ccc} \|\mathbb{E}\mathcal{G}\| & \xrightarrow{f} & \mathcal{G}_0 \\ \downarrow & & \downarrow \\ \|\mathcal{G}\| & \xrightarrow{\varphi} & [\mathcal{G}] \end{array}$$

with f a universal weak equivalence. Now, the stackification 2-functor $St_{\mathcal{CG}}$ commutes with finite weak limits, hence, the following is also a 2-Cartesian diagram:

$$\begin{array}{ccc} \|\mathbb{E}\mathcal{G}\| & \xrightarrow{f} & \mathcal{G}_0 \\ \downarrow & & \downarrow \\ \|\mathcal{G}\| & \xrightarrow{\bar{\varphi}} & \mathcal{X}. \end{array}$$

Since the map $\mathcal{G}_0 \rightarrow \mathcal{X}$ is a $\mathcal{CG}$ -epimorphism and f is a universal weak equivalence, it follows that $\bar{\varphi}$ is a universal weak equivalence. $\square$

We have the following two Corollaries:

Corollary 4.4. *The weak homotopy type of a compactly generated stack is the weak homotopy type of any topological groupoid presenting it.*

Corollary 4.5. *Let $\phi : \mathcal{G} \rightarrow \mathcal{H}$ be a $\mathcal{CG}$ -Morita equivalence between two topological groupoids $\mathcal{G}$ and $\mathcal{H}$. Then ϕ induces a weak homotopy equivalence*

$$\|\phi\| : \|\mathcal{G}\| \rightarrow \|\mathcal{H}\|.$$

Proof. Let $\mathcal{X} := [\mathcal{G}]_{\mathcal{CG}} \cong [\mathcal{H}]_{\mathcal{CG}}$. Then each atlas $\|\mathcal{G}\| \rightarrow \mathcal{X}$ and $\|\mathcal{H}\| \rightarrow \mathcal{X}$ is a universal weak equivalence. The following diagram 2-commutes (with the outer square Cartesian):

$$\begin{array}{ccc}
\|\mathcal{G}\| \times_{\mathcal{X}} \|\mathcal{H}\| & \xrightarrow{\beta} & \|\mathcal{H}\| \\
\alpha \downarrow & \nearrow \|\varphi\| & \downarrow \\
\|\mathcal{G}\| & \xrightarrow{\quad} & \mathcal{X}
\end{array}$$

Since each atlas is a universal weak equivalence, α and β are weak equivalences, and hence so is $\|\varphi\|$. $\square$

4.4. Comparison with Topological Stacks. First, we make a basic observation:

Proposition 4.2. *Let $\mathcal{X}$ be a topological stack. Then the unit map*

$$\varsigma_{\mathcal{X}} : \mathcal{X} \rightarrow St_{\mathcal{CG}}(\mathcal{X})$$

induces an equivalence of groupoids

$$\mathcal{X}(Y) \rightarrow St_{\mathcal{CG}}\mathcal{X}(Y)$$

for all locally compact Hausdorff spaces Y . Moreover, ς is a universal weak equivalence.

Proof. The first statement follows immediately from Proposition 3.5. Suppose $\mathcal{G}$ is a topological groupoid for which $\mathcal{X} \cong [\mathcal{G}]$. Then the $\mathcal{CG}$ -atlas for $St_{\mathcal{CG}}(\mathcal{X})$ given by

$$\|\mathcal{G}\| \rightarrow St_{\mathcal{CG}}(\mathcal{X})$$

factors as

$$\|\mathcal{G}\| \rightarrow \mathcal{X} \xrightarrow{\varsigma_{\mathcal{X}}} St_{\mathcal{CG}}(\mathcal{X}).$$

Since the both

$$\|\mathcal{G}\| \rightarrow \mathcal{X}$$

and

$$\|\mathcal{G}\| \rightarrow St_{\mathcal{CG}}(\mathcal{X})$$

are universal weak equivalences, it follows that so is $\varsigma_{\mathcal{X}}$. $\square$

In particular, to any topological stack, there is a canonically associated compactly generated stack of the same weak homotopy type which restricts to the same stack over locally compact Hausdorff spaces. Conversely, if $\mathcal{Y} \cong [\mathcal{H}]_{\mathcal{CG}}$ is a compactly generated stack, $[\mathcal{H}]$ is an associated topological stack for which the same is true.

Proposition 4.3. *Suppose that $\mathcal{X}$ and $\mathcal{Y}$ are topological stacks and $\mathcal{Y} \cong [\mathcal{H}]$ with $\mathcal{H}_0$ and $\mathcal{H}_1$ compact Hausdorff. Then $\text{Map}(\mathcal{Y}, \mathcal{X})$ is a topological stack, $\text{Map}(St_{\mathcal{CG}}(\mathcal{Y}), St_{\mathcal{CG}}(\mathcal{X}))$ is a compactly generated stack, and they both restrict to the same stack over locally compact Hausdorff spaces. Moreover, the canonical map*

$$\text{Map}(\mathcal{Y}, \mathcal{X}) \rightarrow \text{Map}(St_{\mathcal{CG}}(\mathcal{Y}), St_{\mathcal{CG}}(\mathcal{X}))$$

is a universal weak equivalence.

Proof. In the proof of Lemma 4.6, we show that if $\text{Map}(\mathcal{Y}, \mathcal{X}) \cong [\mathcal{K}]$, then $\text{Map}(St_{\mathcal{CG}}(\mathcal{Y}), St_{\mathcal{CG}}(\mathcal{X})) \cong [\mathcal{K}]_{\mathcal{CG}}$. The result now follows from Proposition 4.2. $\square$

Theorem 4.10. *Let $\mathcal{X}$ and $\mathcal{Y}$ be topological stacks such that $\mathcal{Y} \cong [\mathcal{H}]$ with $\mathcal{H}_0$ and $\mathcal{H}_1$ locally compact. Then $\mathrm{Map}(\mathcal{Y}, \mathcal{X})$ is a paratopological stack, $\mathrm{Map}(St_{\mathcal{CG}}(\mathcal{Y}), St_{\mathcal{CG}}(\mathcal{X}))$ is a compactly generated stack, and there is a canonical universal weak equivalence*

$$\mathrm{Map}(\mathcal{Y}, \mathcal{X}) \rightarrow \mathrm{Map}(St_{\mathcal{CG}}(\mathcal{Y}), St_{\mathcal{CG}}(\mathcal{X})).$$

Moreover, $\mathrm{Map}(\mathcal{Y}, \mathcal{X})$ and $\mathrm{Map}(St_{\mathcal{CG}}(\mathcal{Y}), St_{\mathcal{CG}}(\mathcal{X}))$ restrict to the same stack over locally compact paracompact Hausdorff spaces.

Proof. The unit map $\varsigma_{\mathcal{X}} : \mathcal{X} \rightarrow St_{\mathcal{CG}}(\mathcal{X})$ induces a map

$$\mathrm{Map}(\mathcal{Y}, \mathcal{X}) \rightarrow \mathrm{Map}(\mathcal{Y}, St_{\mathcal{CG}}(\mathcal{X}))$$

by composition. Note that since we have the 2-adjunction $St_{\mathcal{CG}} \dashv \ell$, for any space T :

$$\begin{aligned} \mathrm{Map}(\mathcal{Y}, St_{\mathcal{CG}}(\mathcal{X}))(T) &\cong \mathrm{Hom}_{Gpd^{CGHP}}(\mathcal{Y} \times T, St_{\mathcal{CG}}(\mathcal{X})) \\ &\cong \mathrm{Hom}_{Gpd^{CGHP}}(St_{\mathcal{CG}}(\mathcal{Y}) \times T, St_{\mathcal{CG}}(\mathcal{X})) \\ &\cong \mathrm{Map}(St_{\mathcal{CG}}(\mathcal{Y}), St_{\mathcal{CG}}(\mathcal{X}))(T). \end{aligned}$$

Denote by r the induced map $\mathrm{Map}(\mathcal{Y}, \mathcal{X}) \rightarrow \mathrm{Map}(St_{\mathcal{CG}}(\mathcal{Y}), St_{\mathcal{CG}}(\mathcal{X}))$.

By Theorem 2.7, $\mathrm{Map}(\mathcal{Y}, \mathcal{X})$ is paratopological. Hence there exists a topological stack $[\mathcal{K}]$ and a map

$$q : [\mathcal{K}] \rightarrow \mathrm{Map}(\mathcal{Y}, \mathcal{X})$$

which induces an equivalence of groupoids

$$q(Z) : [\mathcal{K}](Z) \rightarrow \mathrm{Map}(\mathcal{Y}, \mathcal{X})(Z)$$

for every paracompact space Z .

Let T be a compact Hausdorff space:

$$\begin{aligned} \mathrm{Hom}_{Gpd^{CGHP}}(T, [\mathcal{K}]_{\mathcal{CG}}) &\cong \mathrm{Hom}_{Gpd^{CGHP}}(T, [\mathcal{K}]) \\ &\cong \mathrm{Map}(\mathcal{Y}, \mathcal{X})(T) \\ &\cong \mathrm{Hom}_{Gpd^{CGHP}}(\mathcal{Y} \times T, \mathcal{X}) \\ &\cong \mathrm{Hom}_{Gpd^{CGHP}}(St_{\mathcal{CG}}(\mathcal{Y}) \times T, St_{\mathcal{CG}}(\mathcal{X})) \\ &\cong \mathrm{Map}(St_{\mathcal{CG}}(\mathcal{Y}), St_{\mathcal{CG}}(\mathcal{X}))(T), \end{aligned}$$

by virtue of Corollary 4.1. Since $\mathrm{Map}(St_{\mathcal{CG}}(\mathcal{Y}), St_{\mathcal{CG}}(\mathcal{X}))$ is a $\mathcal{CG}$ -stack, it is determined by its restriction to $\mathbb{CH}$, hence

$$(2) \quad \mathrm{Map}(St_{\mathcal{CG}}(\mathcal{Y}), St_{\mathcal{CG}}(\mathcal{X})) \cong [\mathcal{K}]_{\mathcal{CG}}.$$

Therefore, the following diagram 2-commutes:

$$\begin{array}{ccc}
[\mathcal{K}] & \xrightarrow{q} & \mathbb{M}\mathrm{ap}(\mathcal{X}, \mathcal{Y}) \\
\varsigma_{[\mathcal{K}]} \downarrow & \swarrow r & \\
[\mathcal{K}]_{\mathcal{CG}} & &
\end{array}$$

From Proposition 4.2, $\varsigma_{[\mathcal{K}]}$ is a universal weak equivalence. Furthermore from Proposition 2.3, it follows that q is as well. Hence r is a universal equivalence too, as desired.

Finally, since $\mathbb{M}\mathrm{ap}(\mathcal{Y}, \mathcal{X})$ is paratopological, it agrees with $[\mathcal{K}]$ on all paracompact spaces, and since $\mathbb{M}\mathrm{ap}(St_{\mathcal{CG}}(\mathcal{Y}), St_{\mathcal{CG}}(\mathcal{X}))$ is a compactly generated stack, in light of (2), it agrees with $[\mathcal{K}]$ on locally compact Hausdorff spaces. Hence $\mathbb{M}\mathrm{ap}(\mathcal{Y}, \mathcal{X})$ and $\mathbb{M}\mathrm{ap}(St_{\mathcal{CG}}(\mathcal{Y}), St_{\mathcal{CG}}(\mathcal{X}))$ restrict to the same stack on locally compact paracompact Hausdorff spaces. $\square$

We will now give a concrete description of the mapping stack of two compactly generated stacks. Note that since the inclusion of all compact subsets of a space is a $\mathcal{CG}$ -cover, every compactly generated stack has a paracompact Hausdorff atlas (which is also locally compact Hausdorff).

Theorem 4.11. *Let $\mathcal{X} \cong [\mathcal{G}]_{\mathcal{CG}}$ and $\mathcal{Y} \cong [\mathcal{H}]_{\mathcal{CG}}$ be two compactly generated stacks. Assume (without loss of generality) that $\mathcal{H}_0$ is paracompact Hausdorff. Then*

$$[Fib(\mathcal{G})^{\mathcal{H}}]_{\mathcal{CG}} \cong \mathbb{M}\mathrm{ap}(\mathcal{Y}, \mathcal{X}).$$

Proof. It suffices to check that $[Fib(\mathcal{G})^{\mathcal{H}}]_{\mathcal{CG}}$ and $\mathbb{M}\mathrm{ap}(\mathcal{Y}, \mathcal{X})$ agree on every compact Hausdorff space T . Following the same proof as Theorem 2.8, one only has to realize that the product of a compact Hausdorff space with a paracompact space is paracompact. The rest of the proof is identical. $\square$

Corollary 4.6. *If $\mathcal{X}$ and $\mathcal{Y}$ are topological stacks, such that $\mathcal{Y}$ admits an atlas by a CW-complex, then there is a canonical universal weak equivalence*

$$\mathbb{M}\mathrm{ap}(\mathcal{Y}, \mathcal{X}) \rightarrow \mathbb{M}\mathrm{ap}(St_{\mathcal{CG}}(\mathcal{Y}), St_{\mathcal{CG}}(\mathcal{X})),$$

Moreover, $\mathbb{M}\mathrm{ap}(\mathcal{Y}, \mathcal{X})$ and $\mathbb{M}\mathrm{ap}(St_{\mathcal{CG}}(\mathcal{Y}), St_{\mathcal{CG}}(\mathcal{X}))$ restrict to the same stack over locally compact paracompact Hausdorff spaces.

APPENDIX A. STACKS

Let $(\mathcal{C}, J)$ be a small Grothendieck site and let Gpd denote the 2-category of groupoids⁴. Denote by $Gpd^{\mathcal{C}^{op}}$ the 2-category of weak presheaves $\mathcal{C}^{op} \rightarrow Gpd$. That is, weak contravariant 2-functors from $\mathcal{C}$ to Gpd . We call this the 2-category of **prestacks** on $\mathcal{C}$. There exists a canonical inclusion

$$(\cdot)^{id} : \mathrm{Set}^{\mathcal{C}^{op}} \rightarrow Gpd^{\mathcal{C}^{op}},$$

where, each presheaf F is sent to the prestack which assigns to each object C the category $(F(C))^{id}$ whose objects are $F(C)$ and whose arrows are all identities. If C is an object of $\mathcal{C}$, we usually denote $(y_C)^{id}$ simply by C .

⁴Technically, we mean those groupoids which are equivalent to a small category.

Remark. This notion of prestack is non-standard. Typically, the name prestack is reserved for those weak presheaves of groupoids which are separated (see Definition A.1.) See for instance [5]. We choose to call all weak presheaves of groupoids prestacks to emphasize the analogy with presheaves of sets.

We also have a version of the Yoneda lemma:

Lemma A.1. [5] *The 2-Yoneda Lemma: If C is an object of $\mathcal{C}$ and $\mathcal{X}$ a prestack, then there is a natural equivalence of groupoids*

$$\mathrm{Hom}_{Gpd^{\mathcal{C}^{op}}}(C, \mathcal{X}) \cong \mathcal{X}(C).$$

Remark. The 2-category $Gpd^{\mathcal{C}^{op}}$ is both complete and cocomplete; weak limits are computed “pointwise”:

$$\left(\varprojlim \mathcal{X}_i \right) (X) = \varprojlim \mathcal{X}_i (X)$$

where the weak limit to the right is computed in Gpd . Similarly for weak colimits.

Definition A.1. A prestack $\mathcal{X}$ is called a **stack** if for every object C and covering sieve S , then natural map

$$\mathrm{Hom}_{Gpd^{\mathcal{C}^{op}}}(C, \mathcal{X}) \rightarrow \mathrm{Hom}_{Gpd^{\mathcal{C}^{op}}}(S, \mathcal{X})$$

is an equivalence of groupoids.

If this map is only fully faithful, $\mathcal{X}$ is called **separated**.

We denote the full sub-2-category of $Gpd^{\mathcal{C}^{op}}$ consisting of those prestacks that are stacks by $St_J(\mathcal{C})$.

It is immediate from the definition that the weak limit of any small diagram of stacks is again a stack.

If $\mathcal{B}$ is a basis for the topology J , then it suffices to check this condition for every sieve of the form $S_{\mathcal{U}}$, where $\mathcal{U}$ is a covering family. Namely, a prestack is a stack if and only if for every covering family $\mathcal{U} = (U_i \rightarrow C)_i$ the induced map

$$\mathcal{X}(C) \rightarrow \varprojlim \left[\prod \mathcal{X}(U_i) \rightrightarrows \prod \mathcal{X}(U_{ij}) \rightrightarrows \prod \mathcal{X}(U_{ijk}) \right]$$

is an equivalence of groupoids. $\mathcal{X}$ is separated if and only if this map is fully faithful.

The associated groupoid

$$\mathfrak{Des}(\mathcal{X}, \mathcal{U}) := \varprojlim \left[\prod \mathcal{X}(U_i) \rightrightarrows \prod \mathcal{X}(U_{ij}) \rightrightarrows \prod \mathcal{X}(U_{ijk}) \right],$$

obtained as weak limit of the above diagram of groupoids, is called the category of **descent data** for $\mathcal{X}$ at $\mathcal{U}$.

Proposition A.1. *If J is subcanonical and $(U_i \rightarrow C)_i$ is a covering family for an object C , then, in the 2-category of stacks*

$$C \cong \varinjlim \left[\prod U_{ijk} \rightrightarrows \prod U_{ij} \rightrightarrows \prod U_i \right]$$

We will often simply write

$$C \cong \varinjlim_{U_i \rightarrow C} U_i.$$

Definition A.2. A morphism $\varphi : \mathcal{X} \rightarrow \mathcal{Y}$ of stacks is called **representable** if for any object $C \in \mathcal{C}$ and any morphism $C \rightarrow \mathcal{Y}$, the weak pullback $C \times_{\mathcal{Y}} \mathcal{X}$ in the category of stacks is (equivalent to) an object D of $\mathcal{C}$.

We can now define a 2-functor

$$(\cdot)^+ : \mathbf{Gpd}^{\mathcal{C}^{op}} \rightarrow \mathbf{Gpd}^{\mathcal{C}^{op}}$$

by

$$\mathcal{X}^+(C) := \varinjlim_{S \in \mathbf{Cov}(C)} \mathrm{Hom}_{\mathbf{Gpd}^{\mathcal{C}^{op}}}(S, F).$$

We can alternatively define $(\cdot)^+$ by the equation

$$\mathcal{X}^+(C) := \varinjlim_{\mathcal{U} \in \mathbf{cov}(C)} \mathfrak{Des}(\mathcal{X}, \mathcal{U})$$

and obtain a naturally isomorphic 2-functor.

Remark. The weak colimit in either definition must be indexed over a suitable 2-category of covers.

$\mathcal{X}$ is separated if and only if the canonical map $\mathcal{X} \rightarrow \mathcal{X}^+$ is a fully faithful. $\mathcal{X}$ is a stack if and only if this map is an equivalence.

If $\mathcal{X}$ is separated, then $\mathcal{X}^+$ is a stack. Furthermore, $\mathcal{X}^+$ is always separated, for any $\mathcal{X}$. Hence, $\mathcal{X}^{++}$ is always a stack.

Definition A.3. We denote by St_J the 2-functor $\mathcal{X} \mapsto \mathcal{X}^{++}$. It is called the **stackification** 2-functor.

If $\mathcal{X}$ is separated, $St_J(\mathcal{X}) \cong \mathcal{X}^+$, and if $\mathcal{X}$ is a stack, $St_J(\mathcal{X}) \cong \mathcal{X}$. St_J is left-2-adjoint to the inclusion

$$i : St_J(\mathcal{C}) \hookrightarrow \mathbf{Gpd}^{\mathcal{C}^{op}}$$

and preserves finite weak limits.

Remark. $St_J(\mathcal{C})$ is both complete and cocomplete. Since i is a right adjoint, it follows that the computation of weak limits of stacks can be done in the category $\mathbf{Gpd}^{\mathcal{C}^{op}}$, hence can be done “pointwise”. To compute the weak colimit of a diagram of stacks, one must first compute it in $\mathbf{Gpd}^{\mathcal{C}^{op}}$ and then apply the stackification functor St_J .

We end by remarking that the 2-category $St_J(\mathcal{C})$ is Cartesian closed. The exponent $\mathcal{X}^{\mathcal{Y}}$ of two stacks is given by

$$\mathcal{X}^{\mathcal{Y}}(C) = \mathrm{Hom}_{\mathbf{Gpd}^{\mathcal{C}^{op}}}(\mathcal{Y} \times C, \mathcal{X}),$$

and satisfies

$$\mathrm{Hom}_{\mathbf{Gpd}^{\mathcal{C}^{op}}}(\mathcal{Z}, \mathcal{X}^{\mathcal{Y}}) \cong \mathrm{Hom}_{\mathbf{Gpd}^{\mathcal{C}^{op}}}(\mathcal{Y} \times \mathcal{Z}, \mathcal{X})$$

for all stacks $\mathcal{Z}$.

APPENDIX B. TOPOLOGICAL GROUPOIDS AND TOPOLOGICAL STACKS

B.1. Definitions and Some Examples. This section is a review of topological groupoids and topological stacks. The material is entirely standard aside from the fact that some of the standard results are stated for more general Grothendieck topologies.

Definition B.1. A **topological groupoid** is a groupoid object in $\mathbf{TOP}$. Explicitly, it is a diagram

$$\begin{array}{ccc} \mathcal{G}_1 \times_{\mathcal{G}_0} \mathcal{G}_1 & \xrightarrow{m} & \mathcal{G}_1 \\ & \begin{array}{c} \text{\scriptsize i} \curvearrowright \\ \text{\scriptsize s} \rightarrow \\ \text{\scriptsize t} \rightarrow \\ \text{\scriptsize u} \curvearrowleft \end{array} & \mathcal{G}_0 \end{array}$$

of topological spaces and continuous maps satisfying the usual axioms. Forgetting the topological structure (i.e. applying the forgetful functor from $\mathbf{TOP}$ to $\mathbf{Set}$), one obtains an ordinary (small) groupoid.

Topological groupoids form a 2-category with continuous functors as 1-morphisms and continuous natural transformations as 2-morphisms. We denote the 2-category of topological groupoids by $\mathbf{TOPGpd}$.

Remark. $\mathbf{TOPGpd}$ has weak fibred products and arbitrary products. Hence, it follows from [25] and [6] that it is a complete 2-category.

Definition B.2. Given a topological space X , we denote by $(X)^{id}$ the topological groupoid whose object and arrow space are both X and all of whose structure maps are the identity morphism of X . The arrow space is the collection of all the identity arrows for the objects X .

Definition B.3. Given a topological space X , the **pair groupoid** $Pair(X)$ is the topological groupoid whose object space is X and whose arrow space is $X \times X$, where an element

$$(x, y) \in X \times X$$

is viewed as an arrow from y to x and composition is defined by the rule

$$(x, y) \cdot (y, z) = (x, z).$$

Definition B.4. Given a continuous map $\phi : U \rightarrow X$, the **relative pair groupoid** $Pair(\phi)$ is defined to be the topological groupoid whose arrow space is the fibred product $U \times_X U$ and object space is U , where an element

$$(x, y) \in U \times_X U \subset U \times U$$

is viewed as an arrow from y to x and composition is defined by the rule

$$(x, y) \cdot (y, z) = (x, z).$$

The pair groupoid of a space X is the relative pair groupoid of the unique map from X to the one-point space.

Definition B.5. Given a topological groupoid $\mathcal{G}$ and a continuous map $f : X \rightarrow \mathcal{G}_0$, there is an **induced groupoid** $f^*(\mathcal{G})$, which is a topological groupoid whose object space is X such that arrows between x and y in $f^*(\mathcal{G})$ are in bijection with arrows between $f(x)$ and $f(y)$ in $\mathcal{G}$. When $X = \coprod_{\alpha} U_{\alpha}$ with $\mathcal{U} = (U_{\alpha} \hookrightarrow X)_{\alpha}$ an open cover

of $\mathcal{G}_0$ and $X \rightarrow \mathcal{G}_0$ the canonical map, $f^*(\mathcal{G})$ is denoted by $\mathcal{G}_{\mathcal{U}}$. If in addition to this, $\mathcal{G} = (T)^{id}$ for some topological space T , then this is called the **Čech groupoid** associated to the cover $\mathcal{U}$ of T and is denoted by $T_{\mathcal{U}}$.

Remark. If the open cover $\mathcal{U}$ is instead a cover for a different Grothendieck topology on $\mathbf{TOP}$, the above still makes sense. This will be important later.

B.2. Principal Bundles. Principal bundles for topological groups, and more generally for topological groupoids, are classical objects of study. However, principal bundles (and many other objects involving a local triviality condition) should not be thought of as objects associated to the *category* $\mathbf{TOP}$, but rather as objects associated to the *Grothendieck site* $(\mathbf{TOP}, \mathcal{O})$, where $\mathcal{O}$ is the **open cover topology** on $\mathbf{TOP}$. This Grothendieck topology is defined by declaring a family of maps $(O_{\alpha} \rightarrow X)_{\alpha}$ to be a covering family if and only if it constitutes an open cover of X . The concept of principal bundles generalizes to other Grothendieck topologies on $\mathbf{TOP}$ and we will need this generality later when we introduce the compactly generated Grothendieck topology. For the remainder of this subsection, let J be an arbitrary subcanonical Grothendieck topology on $\mathbf{TOP}$.

Definition B.6. Given a topological groupoid $\mathcal{G}$, a (left) $\mathcal{G}$ -**space** is a space E equipped with a **moment map** $\mu : E \rightarrow \mathcal{G}_0$ and an **action map** $\rho : \mathcal{G}_1 \times_{\mathcal{G}_0} E \rightarrow E$, where

$$\begin{array}{ccc} \mathcal{G}_1 \times_{\mathcal{G}_0} E & \longrightarrow & E \\ \downarrow & & \downarrow \mu \\ \mathcal{G}_1 & \xrightarrow{s} & \mathcal{G}_0 \end{array}$$

is the fibred product, such that the following conditions hold:

- i) $(gh) \cdot e = g \cdot (h \cdot e)$ whenever e is an element of E and g and h elements of $\mathcal{G}_1$ with domains such that the composition makes sense
- ii) $1_{\mu(e)} \cdot e = e$ for all $e \in E$
- iii) $\mu(g \cdot e) = t(g)$ for all $g \in \mathcal{G}_1$ and $e \in E$.

Definition B.7. Suppose that $\mathcal{G} \mathfrak{C} E$ is a $\mathcal{G}$ -space. Then the **action groupoid** $\mathcal{G} \ltimes E$ is defined to be the topological groupoid whose arrow space is $\mathcal{G} \times_{\mathcal{G}_0} E$ and whose object space is E . An element

$$(g, e) \in \mathcal{G} \times_{\mathcal{G}_0} E \subset \mathcal{G} \times E$$

is viewed as an arrow from e to $g \cdot e$. Composition is defined by the rule

$$(g, e) \cdot (h, g \cdot e) = (hg, e).$$

Definition B.8. Given a topological groupoid $\mathcal{G}$, the **translation groupoid** $\mathbb{E}\mathcal{G}$ is defined to be the action groupoid $\mathcal{G} \ltimes \mathcal{G}_1$ with respect to the action of $\mathcal{G}$ on $\mathcal{G}_1$ by multiplication.

Definition B.9. A (left) $\mathcal{G}$ -**bundle** over a space X (with respect to J) is a (left) $\mathcal{G}$ -space P equipped with $\mathcal{G}$ -invariant **projection map**

$$\pi : P \rightarrow X$$

which **admits local sections** with respect to the Grothendieck topology J . This last condition means that there exists a covering family $\mathcal{U} = (U_i \rightarrow X)_i$ in J and

morphisms $\sigma_i : U_i \rightarrow P$ called **local sections** such that the following diagram commutes for all i :

$$\begin{array}{ccc} & P & \\ \sigma_i \nearrow & & \searrow f \\ U_i & \xrightarrow{\quad} & X. \end{array}$$

This condition is equivalent to requiring that the projection map is a J -cover. Such a $\mathcal{G}$ -bundle is called **(J)-principal** if the induced map,

$$\mathcal{G}_1 \times_{\mathcal{G}_0} P \rightarrow P \times_X P$$

is a homeomorphism.

We typically denote such a principal bundle as

$$\begin{array}{ccc} \mathcal{G}_1 & \xrightarrow{\quad \mathcal{C} \quad} & P \\ \downarrow & \mu \nearrow & \downarrow \pi \\ \mathcal{G}_0 & & X. \end{array}$$

Remark. To ease terminology, for the rest of this section, the term principal bundle, will refer to a J -principal bundle for our fixed topology J .

Definition B.10. Any topological groupoid $\mathcal{G}$ determines a principal $\mathcal{G}$ -bundle over $\mathcal{G}_0$ by

$$\begin{array}{ccc} \mathcal{G}_1 & \xrightarrow{\quad \mathcal{C} \quad} & \mathcal{G}_1 \\ \downarrow & t \nearrow & \downarrow s \\ \mathcal{G}_0 & & \mathcal{G}_0 \end{array}$$

called the **unit bundle**, $1_{\mathcal{G}}$.

Definition B.11. Given P and P' , two principal $\mathcal{G}$ -bundle over X , a continuous map $f : P \rightarrow P'$ is a **map of principal bundles** if it respects the projection maps and is $\mathcal{G}$ -equivariant. It is easy to check that any such map must be an isomorphism of principal bundles.

Definition B.12. Let $f : Y \rightarrow X$ be a map and suppose that P is a principal $\mathcal{G}$ -bundle over X . Then we can give $Y \times_X P \rightarrow Y$ the structure of a principal $\mathcal{G}$ -bundle $f^*(P)$ over Y , called the **pull-back bundle**, in the obvious way.

Definition B.13. If $\mathcal{G}$ is topological groupoid and P is a principal $\mathcal{G}$ -bundle over X , then we define the **gauge groupoid** $\text{Gauge}(P)$ to be the following topological groupoid:

The fibred product

$$\begin{array}{ccc} P \times_{\mathcal{G}_0} P & \xrightarrow{\quad} & P \\ \downarrow & & \downarrow \mu \\ P & \xrightarrow{\quad \mu \quad} & \mathcal{G}_0 \end{array}$$

admits a left- $\mathcal{G}$ -action with moment map $\tilde{\mu}((p, q)) = \mu(p)$ via

$$g \cdot (p, q) = (g \cdot p, g \cdot q).$$

The arrow space of $Gauge(P)$ is the quotient $P \times_{\mathcal{G}_0} P / \mathcal{G}$ and the object space is X . An equivalence class $[(p, q)]$ is viewed as an arrow from $\pi(p)$ to $\pi(q)$ (which is well defined as π is $\mathcal{G}$ -invariant.) Composition is determined by the rule

$$[(p, q)] \cdot [(q', r)] = [(p, g \cdot r)]$$

where g is the unique element of $\mathcal{G}_1$ such that $g \cdot q' = q$.

Definition B.14. Let $\mathcal{G}$ and $\mathcal{H}$ be topological groupoids. A (left) **principal $\mathcal{G}$ -bundle over $\mathcal{H}$** is a (left) principal $\mathcal{G}$ -bundle

$$\begin{array}{ccc} \mathcal{G}_1 & \xrightarrow{\mathcal{C}} & P \\ \downarrow & \searrow \mu & \downarrow \nu \\ \mathcal{G}_0 & & \mathcal{H}_0 \end{array}$$

over $\mathcal{H}_0$, such that P also has the structure of a right $\mathcal{H}$ -bundle with moment map ν , with the $\mathcal{G}$ and $\mathcal{H}$ actions commuting in the obvious sense. We typically denote such a bundle by

$$\begin{array}{ccccc} \mathcal{G}_1 & \xrightarrow{\mathcal{C}} & P & \xrightarrow{\mathcal{H}} & \mathcal{H}_1 \\ \downarrow & \searrow \mu & & \searrow \nu & \downarrow \\ \mathcal{G}_0 & & & & \mathcal{H}_0 \end{array}$$

Remark. To a continuous functor $\varphi : \mathcal{H} \rightarrow \mathcal{G}$, one can canonically associate a principal $\mathcal{G}$ -bundle over $\mathcal{H}$. It is defined by putting the obvious (right) $\mathcal{H}$ -bundle structure on the total space of the pullback bundle $\varphi_0^*(1_{\mathcal{G}}) = \mathcal{G}_1 \times_{\mathcal{G}_0} \mathcal{H}_0$.

B.3. Bicategories of Topological Groupoids.

Definition B.15. A continuous functor $\varphi : \mathcal{H} \rightarrow \mathcal{G}$ between two topological groupoids is a **J -Morita equivalence** if the following two properties hold:

i) (Essentially Surjective)

The map $t \circ pr_1 : \mathcal{G}_1 \times_{\mathcal{G}_0} \mathcal{H}_0 \rightarrow \mathcal{G}_0$ admits local sections with respect to the topology J , where $\mathcal{G}_1 \times_{\mathcal{G}_0} \mathcal{H}_0$ is the fibred product

$$\begin{array}{ccc} \mathcal{G}_1 \times_{\mathcal{G}_0} \mathcal{H}_0 & \xrightarrow{pr_2} & \mathcal{H}_0 \\ pr_1 \downarrow & & \downarrow \varphi \\ \mathcal{G}_1 & \xrightarrow{s} & \mathcal{G}_0. \end{array}$$

i) (Fully Faithful)

The following is a fibred product

$$\begin{array}{ccc} \mathcal{H}_1 & \xrightarrow{\varphi} & \mathcal{G}_1 \\ \langle s, t \rangle \downarrow & & \downarrow \langle s, t \rangle \\ \mathcal{H}_0 \times \mathcal{H}_0 & \xrightarrow{\varphi \times \varphi} & \mathcal{G}_0 \times \mathcal{G}_0. \end{array}$$

Remark. If $\mathcal{U}$ is a J -cover of the object space $\mathcal{G}_0$ of a topological groupoid $\mathcal{G}$, then the induced map $\mathcal{G}_{\mathcal{U}} \rightarrow \mathcal{G}$ is a J -Morita equivalence.

Remark. We will again suppress the reference to the Grothendieck topology J ; for the rest of the section a Morita equivalence will implicitly mean a J -Morita equivalence for our fixed topology J . A Morita equivalence with respect to the open cover topology will be called an ordinary Morita equivalence.

Remark. The property of being a Morita equivalence is weaker than being an equivalence in the 2-category $\mathbf{TOPGpd}$. In fact, a Morita equivalence is an equivalence in $\mathbf{TOPGpd}$ if and only if $t \circ pr_1$ admits a global section. However, any Morita equivalence does induce an equivalence in the 2-category $\mathbf{Gpd}$ after applying the forgetful 2-functor. Morita equivalences are sometimes referred to as weak equivalences.

We denote by W_J the class of Morita equivalences. The class W_J admits a right calculus of fractions in the sense of [22]. There is a bicategory $\mathbf{TOPGpd}[W_J^{-1}]$ obtained from the 2-category $\mathbf{TOPGpd}$ by formally inverting the Morita equivalences. A 1-morphism from $\mathcal{H}$ to $\mathcal{G}$ in this bicategory is a diagram of continuous functors

$$\begin{array}{ccc} & \mathcal{K} & \\ w \swarrow & & \searrow \\ \mathcal{H} & & \mathcal{G} \end{array}$$

such that w is a Morita equivalence. Such a diagram is called a **generalized homomorphism**. There is also a well defined notion of a 2-morphism. For details see [22].

There is in fact another bicategory which one can construct from the 2-category $\mathbf{TOPGpd}$. This bicategory, denoted by $\mathbf{Bun}^J\mathbf{TOPGpd}$ again has the same objects as $\mathbf{TOPGpd}$. A 1-morphism between two topological groupoids $\mathcal{H}$ and $\mathcal{G}$ is a left-principal $\mathcal{G}$ -bundle over $\mathcal{H}$ (with respect to J). There is a well-defined way of composing principal bundles, for details see [15] or [16]. A 2-morphism between two such principal bundles is a biequivariant map (a map which is equivariant with respect to the left- $\mathcal{G}$ -action and the right- $\mathcal{H}$ -action). A principal $\mathcal{G}$ -bundle over $\mathcal{H}$ is an equivalence in this bicategory if and only if the underlying $\mathcal{H}$ -bundle is also principal.

By the remark following definition B.14, there is a canonical inclusion

$$\mathbf{TOPGpd} \hookrightarrow \mathbf{Bun}^J\mathbf{TOPGpd}$$

which sends Morita equivalences to equivalences. Therefore, there is an induced map

$$\mathbf{TOPGpd}[W_J^{-1}] \rightarrow \mathbf{Bun}^J\mathbf{TOPGpd}$$

of bicategories.

Theorem B.1. *The induced map*

$$\mathbf{TOPGpd}[W_J^{-1}] \rightarrow \mathbf{Bun}^J\mathbf{TOPGpd}$$

is an equivalence of bicategories.

This theorem is well known. A 1-categorical version of this theorem is proven in [15], Section 2.6. The general result follows easily from [22], Section 3.4.

B.4. Topological Stacks. Consider the 2-category $Gpd^{\mathbf{TOP}^{op}}$ of prestacks over $\mathbf{TOP}$ ⁵, that is contravariant (possibly weak) 2-functors from the category $\mathbf{TOP}$ into the 2-category of (essentially small) groupoids Gpd . Let G be a topological group. A standard example would be the prestack that assigns to each space the category of principal G -bundles over that space (this category is a groupoid). More generally, let $\mathcal{G}$ be a topological groupoid. Then $\mathcal{G}$ determines a weak presheaf on $\mathbf{TOP}$ by the rule

$$X \mapsto \mathrm{Hom}_{\mathbf{TOP}Gpd} \left((X)^{(id)}, \mathcal{G} \right).$$

This defines an extended Yoneda 2-functor $\tilde{y} : \mathbf{TOP}Gpd \rightarrow Gpd^{\mathbf{TOP}^{op}}$ and we have the obvious commutative diagram

$$\begin{array}{ccc} \mathbf{TOP} & \xrightarrow{y} & \mathbf{Set}^{\mathbf{TOP}^{op}} \\ (\cdot)^{(id)} \downarrow & & \downarrow (\cdot)^{(id)} \\ \mathbf{TOP}Gpd & \xrightarrow{\tilde{y}} & Gpd^{\mathbf{TOP}^{op}}. \end{array}$$

Remark. $\tilde{y}$ preserves all weak-limits.

Given a subcanonical Grothendieck topology J on $\mathbf{TOP}$, we denote by $[\mathcal{G}]_J$ the associated stack on $(\mathbf{TOP}, J)$, $St_J \circ \tilde{y}\mathcal{G}$, where St_J is the stackification 2-functor (Definition A.3).

It can be checked that

$$[\mathcal{G}]_J(X) \cong \varinjlim_{\mathcal{U} \in Cov(X)} \mathrm{Hom}_{\mathbf{TOP}Gpd}(X_{\mathcal{U}}, \mathcal{G}),$$

where the weak 2-colimit above is taken over a suitable 2-category of J -covers. For details in the case J is the open cover topology, see [7].

Remark. Since St_J preserves finite weak-limits, it follows that $[\cdot]_J$ does as well.

Definition B.16. A stack $\mathcal{X}$ on $(\mathbf{TOP}, J)$ is **presentable** if it is 2-isomorphic to $[\mathcal{G}]_J$ for some topological groupoid $\mathcal{G}$. In this case, $\mathcal{G}$ is said to be a **presentation** of $\mathcal{X}$.

We denote the full sub-2-category of $St_J(\mathbf{TOP})$ consisting of presentable stacks by $\mathfrak{Pres}St_J(\mathbf{TOP})$.

Definition B.17. A **topological stack** is a presentable stack for the open cover topology on $\mathbf{TOP}$. We shall denote the topological stack associated to a topological groupoid $\mathcal{G}$ by $[\mathcal{G}]$. We shall also denote the 2-category of topological stacks by $\mathfrak{TopSt}$.

Theorem B.2. *The 2-functor*

$$St_J \circ \tilde{y} : \mathbf{TOP}Gpd \rightarrow \mathfrak{Pres}St_J(\mathbf{TOP})$$

induces an equivalence of bicategories

⁵Technically, this is not well defined due to set-theoretic issues, however, this can be overcome by careful use of Grothendieck universes. We will not dwell on this and all such similar size issues in this paper.

$$P_J : \text{TOPGpd} [W_J^{-1}] \xrightarrow{\sim} \mathfrak{PresSt}_J(\text{TOP}).$$

This theorem is well known. For example, see [22] for the case of étale topological groupoids and topological stacks with an étale atlas, and also for étale Lie groupoids and étale differentiable stacks. [2] contains much of the necessary ingredients for the proof of the case of general Lie groupoids and differentiable stacks in its so called Dictionary Lemmas. Similar statements in the case of algebraic stacks can be found in [17]. The general theorem follows again from an easy application of [22], Section 3.4.

Hence all three bicategories, $\mathfrak{PresSt}_J(\text{TOP})$, $\text{TOPGpd} [W_J^{-1}]$ and $\text{Bun}^J \text{TOPGpd}$ are equivalent.

Corollary B.1. *If $\mathcal{G}$ is a topological groupoid, then the associated presentable stack $[\mathcal{G}]_J$ (is equivalent to one that) assigns to each topological space T , the groupoid of principal $\mathcal{G}$ -bundles over T , $\text{Bun}_{\mathcal{G}}^J(T)$.*

Definition B.18. [11] A morphism $\varphi : \mathcal{X} \rightarrow \mathcal{Y}$ between prestacks is said to be a **J -covering morphism** if for every space X and every object $x \in \mathcal{Y}(X)_0$, there exists a J -covering family $\mathcal{U} = (f_i : U_i \rightarrow X)_i$ of X such that for each i there exists an object $x_i \in \mathcal{X}(U_i)_0$ and a(n) (iso)morphism

$$\alpha_i : \varphi(\mathcal{X}(f_i)(x_i)) \rightarrow \mathcal{Y}(f_i)(x).$$

If $\mathcal{X}$ and $\mathcal{Y}$ are both J -stacks, J -covering morphisms are referred to as **J -epimorphisms**[11],[18].

There is a more intrinsic description of presentable stacks:

Definition B.19. A **J -atlas** for a stack $\mathcal{X}$ over (TOP, J) is a *representable* (see Definition A.2) J -epimorphism $p : X \rightarrow \mathcal{X}$ from a topological space X .

Proposition B.1. [18] *A stack $\mathcal{X}$ is presentable if and only if it has an atlas.*

Suppose that $p : X \rightarrow \mathcal{X}$ is an atlas. Consider the weak fibred product

$$\begin{array}{ccc} X \times_{\mathcal{X}} X & \longrightarrow & X \\ \downarrow & & \downarrow p \\ X & \xrightarrow{p} & \mathcal{X}. \end{array}$$

Then $X \times_{\mathcal{X}} X \rightrightarrows X$ has the structure of a topological groupoid whose associated stack is $\mathcal{X}$.

Conversely, given a topological groupoid $\mathcal{G}$, the canonical map of groupoids $(\mathcal{G}_0)^{\text{id}} \rightarrow \mathcal{G}$ (which is the identity objects and u on arrows) produces a morphism $p : \mathcal{G}_0 \rightarrow [\mathcal{G}]_J$ which is a representable epimorphism.

We end this section by stating a technical lemma.

Lemma B.3. *The presentable stack $[\mathcal{G}]_J$ is the weak colimit of the following diagram of representables:*

$$\begin{array}{ccccc} \mathcal{G}_1 \times_{\mathcal{G}_0} \mathcal{G}_1 & \rightrightarrows & \mathcal{G}_1 & \xrightarrow{s} & \mathcal{G}_0 \\ & & & \searrow t & \\ & & & \mathcal{G}_0 & \\ & & & \uparrow u & \end{array}$$

where the three parallel arrows are the first and second projections and the composition map.

Proof. This is the content of [18] Proposition 3.19. $\square$

B.5. Representable Maps of Topological Stacks. [18]

Definition B.20. Let P be a property of a map of topological spaces. P is said to be **invariant under change of base** if for all $f : Y \rightarrow X$ with property P , if $g : Z \rightarrow X$ is any continuous map, the induced map $Z \times_X Y \rightarrow Z$ also has property P . P is said to be **invariant under restriction** if this holds whenever g is an embedding. A property P which is invariant under restriction is said to be **local on the target** if any $f : Y \rightarrow X$ for which there exists an open cover $(U_\alpha \rightarrow X)$ such that the induced map $\coprod_\alpha U_\alpha \times_X Y \rightarrow \coprod_\alpha U_\alpha$ has property P , must also have property P .

Examples of such properties are being an open map, covering map, closed map, local homeomorphism etc.

Definition B.21. Let P be a property of a map of topological spaces which is invariant under restriction and local on the target. Then, a representable map (see definition A.2) $f : \mathcal{X} \rightarrow \mathcal{Y}$ of stacks (for the open cover topology) is said to have property P if for any map $T \rightarrow \mathcal{Y}$ from a topological space, the induced map $T \times_{\mathcal{Y}} \mathcal{X} \rightarrow T$ has property P .

Remark. Both of these definitions have obvious modifications to accommodate Grothendieck topologies other than the open cover topology.

REFERENCES

- [1] Alejandro Adem, Johann Leida, and Yongbin Ruan. Orbifolds and stringy topology. *Cambridge Tracts in Mathematics* 171, Cambridge University Press, Cambridge, 2007.
- [2] Kai Behrend, Ping Xu. Differentiable Stacks and Gerbes. *arXiv:math/0605694*
- [3] Weimin Chen. On a notion of maps between orbifolds, I. Function spaces. *Commun. Contemp. Math.* 8 (2006), no. 5, 569620.
- [4] Martín Escardó, Jimmie Lawson, and Alex Simpson. Comparing Cartesian closed categories of (core) compactly generated spaces. *Topology and its Applications* 143 (2004) 105-145
- [5] Barbara Fantechi, Lothar Göttsche, Luc Illusie, Steven Kleiman, Niltin Nitsure, and Angelo Vistoli. *Fundamental Algebraic Geometry: Grothendieck's FGA Explained*, American Mathematical Society, 2006
- [6] Thomas M. Fiore. Pseudo Limits, Biadjoints, and Pseudo Algebras: Categorical Foundations of Conformal Field Theory. *Mem. Amer. Math. Soc.* 182 (2006), no. 860
- [7] David Gepner, André Henriques. Homotopy Theory of Orbispaces. *arXiv:math/0701916v1*
- [8] André Haefliger. On the Space of Morphisms Between Étale Groupoids. *arXiv:0707.4673v2*
- [9] Ernesto Lupercio, and Bernardo Uribe. *Loop groupoids, gerbes, and twisted sectors on orbifolds*, Orbifolds in mathematics and physics (Madison, WI, 2001), 163184, *Contemp. Math.*, 310, Amer. Math. Soc., Providence, RI, 2002.
- [10] Saunders Mac Lane and Ieke Moerdijk, *Sheaves in Geometry and Logic*, Springer-Verlag, New York, NY, 1992.
- [11] David Metzler. Topological and Smooth Stacks. *arXiv:math/0306176v1*
- [12] Ieke Moerdijk. Classifying toposes and foliations. *Ann. Inst. Fourier (Grenoble)* 41:1 (1991), 189209.
- [13] Ieke Moerdijk. On the Weak Homotopy Type of Étale Groupoids, in *Integrable Systems and Foliations*, Birkhäuser (1997), 145-156
- [14] Ieke Moerdijk, and Janez Mrčun, *Introduction to foliations and Lie groupoids*, University Press, Cambridge, 2003

- [15] Ieke Moerdijk, and Janez Mrčun. Lie groupoids, sheaves and cohomology, in *Poisson geometry, deformation quantisation and group representations*. Number 323 in London Math. Soc. Lecture Note Ser. Cambridge Univ. Press, Cambridge, 2005.
- [16] Janez Mrčun. Stability and Invariants of Hilsum-Skandalis Maps. Ph.D. thesis, Utrecht 1996, *arXiv:math/0506484v1 [math.DG]*
- [17] Niko Naumann. The stack of formal groups in stable homotopy theory. *Advances in Math.* 215 (2007)
- [18] Behrang Noohi. Foundations of topological stacks, I. *arXiv:math/0503247v1*
- [19] Behrang Noohi. Homotopy types of topological stacks. *arXiv:math/0808.3799v1*
- [20] Behrang Noohi. Mapping stacks of topological stacks. *arXiv:math/0809.2373v1*
- [21] Ethan Pribble. *A guide to understanding stacks, Part III: Stack theory as lax 2-categorical sheaf theory*, seminar slides, University of Rochester, 2007
- [22] Dorette Pronk. Etendues and stacks as bicategories of fractions. *Compositio Math.*, 102(3):243303, 1996
- [23] Micheal Artin, Alexander Grothendieck, and Jean-Louis Verdier. Théorie des Topos et Cohomologie Étale des Schémas. *Springer Lecture Notes in Math.* 269 (1972)
- [24] Norman E. Steenrod. A convenient category of topological spaces. *Michigan Math. J.*, 14:113152, 1967
- [25] Ross Street. Fibrations in bicategories. *Cahiers de topologie et géométrie différentielle catégoriques*, 21 no. 2 (1980), p. 111-160